

Supplemental Appendix to “A Practical Guide to Compact Infinite Dimensional Parameter Spaces”

Joachim Freyberger* Matthew A. Masten†

July 4, 2018

Abstract

This supplemental appendix provides some formal definitions, useful lemmas, several additional results discussed in the main paper, and proofs for all results in the main paper.

A Some formal definitions and useful lemmas

Let $(X, \|\cdot\|_X)$ and $(Y, \|\cdot\|_Y)$ be normed vector spaces. Then we use the following definitions.

- $A \subseteq X$ is $\|\cdot\|_X$ -*bounded* if there is a scalar $R > 0$ such that $\|x\|_X \leq R$ for all $x \in A$. Equivalently, if A is contained in a $\|\cdot\|_X$ -ball of radius R : $A \subseteq \{x \in X : \|x\|_X \leq R\}$.
- $A \subseteq X$ is $\|\cdot\|_X$ -*relatively compact* if its $\|\cdot\|_X$ -closure is $\|\cdot\|_X$ -compact.
- $(X, \|\cdot\|_X)$ is *embedded* in $(Y, \|\cdot\|_Y)$ if
 1. X is a vector subspace of Y , and
 2. the identity operator $I : X \rightarrow Y$ defined by $Ix = x$ for all $x \in X$ is continuous.

This is also sometimes called being *continuously embedding*, since the identity operator is required to be continuous. Since I is linear, part (2) is equivalent to the existence of a constant M such that

$$\|x\|_Y \leq M\|x\|_X \quad \text{for all } x \in X.$$

Write $X \hookrightarrow Y$ to denote that $(X, \|\cdot\|_X)$ is embedded in $(Y, \|\cdot\|_Y)$.

- $T : X \rightarrow Y$ is a *compact operator* if it maps $\|\cdot\|_X$ -bounded sets to $\|\cdot\|_Y$ -relatively compact sets. That is, $T(A)$ is $\|\cdot\|_Y$ -relatively compact whenever A is $\|\cdot\|_X$ -bounded.
- $(X, \|\cdot\|_X)$ is *compactly embedded* in $(Y, \|\cdot\|_Y)$ if it is embedded and if the identity operator I is compact.

*Department of Economics, University of Wisconsin-Madison, jfreyberger@ssc.wisc.edu

†Department of Economics, Duke University, matt.masten@duke.edu

- A *cone* is a set $C = C(v, a, h, \kappa) = \{v + x \in \mathbb{R}^n : 0 \leq \|x\|_e \leq h, \angle(x, a) \leq \theta\}$. This cone is defined by four parameters: The cone's vertex $v \in \mathbb{R}^n$, an axis direction vector $a \in \mathbb{R}^n$, a height $h \in [0, \infty]$, and an angle parameter $\theta \in (0, 2\pi]$. $\angle(x, a)$ denotes the angle between x and a (let $\angle(x, x) = 0$). $\theta > 0$ ensures that the cone has volume. If $h < \infty$ then we say C is a *finite cone*.
- A set A satisfies the *cone condition* if there is some finite cone C such that for every $x \in A$ the cone C can be moved by rigid motions to have x as its vertex; that is, there is a finite cone C_x with vertex at x which is congruent to C . A sufficient condition for this is that A is a product of intervals, or that A is a ball.

Lemma 1. If all $\|\cdot\|_X$ -balls are $\|\cdot\|_Y$ -relatively compact, then $(X, \|\cdot\|_X)$ is compactly embedded in $(Y, \|\cdot\|_Y)$.

Lemma 1 states that, for proving compact embeddedness, it suffices to show that any $\|\cdot\|_X$ -ball is $\|\cdot\|_Y$ -relatively compact.

Lemma 2. Let $\|\cdot\|_X$ and $\|\cdot\|_Y$ be norms on a vector space A . Suppose A is $\|\cdot\|_X$ -closed and $\|\cdot\|_X \leq C\|\cdot\|_Y$ for $C < \infty$. Then A is $\|\cdot\|_Y$ -closed.

Corollary 1. Let $(\mathcal{F}_j, \|\cdot\|_j)$ be Banach spaces for all $j \in \mathbb{N}$ such that $\mathcal{F}_{j+1} \subseteq \mathcal{F}_j$ and $\|f\|_j \leq C_j \|f\|_{j+1}$ for all $f \in \mathcal{F}_{j+1}$, where $C_j < \infty$. Let

$$\Theta_j = \{f \in \mathcal{F}_j : \|f\|_j \leq C\}.$$

Assume Θ_k is $\|\cdot\|_1$ -closed. Then Θ_k is $\|\cdot\|_j$ -closed for all $1 \leq j < k$.

Lemma 2 says that closedness in a weaker norm can always be converted to closedness in a stronger norm. Lemma 3 gives conditions under which the reverse is true: when we can take closedness in a stronger norm and convert that to closedness in a weaker norm.

Lemma 3. Let $(H_1, \|\cdot\|_1)$ and $(H_2, \|\cdot\|_2)$ be reflexive Banach spaces. Suppose $(H_1, \|\cdot\|_1)$ is compactly embedded in $(H_2, \|\cdot\|_2)$. Let $B < \infty$ be a constant. Then the $\|\cdot\|_1$ -ball

$$\Omega = \{h \in H_1 : \|h\|_1 \leq B\}$$

is $\|\cdot\|_2$ -closed.

Lemma 3 generalizes lemma A.1 of Santos (2012), which assumed both spaces were separable Hilbert spaces. We thank Kengo Kato for pointing out this generalization and its proof. Also recall that all Hilbert spaces are reflexive.

Lemma 4. Let $(X, \|\cdot\|_X)$, $(Y, \|\cdot\|_Y)$, and $(Z, \|\cdot\|_Z)$ be Banach spaces. Suppose

1. $(X, \|\cdot\|_X)$ is compactly embedded in $(Z, \|\cdot\|_Z)$.

2. $(Z, \|\cdot\|_Z)$ is embedded in $(Y, \|\cdot\|_Y)$.

Then $(X, \|\cdot\|_X)$ is compactly embedded in $(Y, \|\cdot\|_Y)$.

Note that assumption 2 implies

$$\{g : \|g\|_Z < \infty\} \subseteq \{g : \|g\|_Y < \infty\}.$$

B Norm inequality lemmas

Lemma 5. Let $\mu : \mathcal{D} \rightarrow \mathbb{R}_+$ be a nonnegative function. Let $m_0, m \geq 0$ be integers. Suppose assumption 4 holds for $\mu = \mu_s$. Then for every compact subset $\mathcal{C} \subseteq \mathcal{D}$, there is a constant $M_{\mathcal{C}} < \infty$ such that

$$\|\mu^{1/2} f\|_{m+m_0, 2, \mathbb{1}_{\mathcal{C}}} \leq M_{\mathcal{C}} \|f\|_{m+m_0, 2, \mu \mathbb{1}_{\mathcal{C}}}$$

for all f such that these norms are defined. If the stronger assumption 3 holds, then this result holds for $\mathcal{C} = \mathcal{D}$ too.

Lemma 5 generalizes lemma A.1 part (a) of Gallant and Nychka (1987) to allow for more general weight functions, as discussed in section 4.1. Note that Gallant and Nychka's (1987) lemma A.1 stated $\sup_{x \in \mathcal{D}} \mu(x) < \infty$ as an additional assumption. This condition is not used in our proof, nor was it used in their proof, which is fortunate since it is violated when μ upweights.

Lemma 6. Let $\mu : \mathcal{D} \rightarrow \mathbb{R}_+$ be a nonnegative function. Let $m \geq 0$ be an integer. Suppose assumption 4 holds for $\mu = \mu_s$. Then for every compact subset $\mathcal{C} \subseteq \mathcal{D}$, there is a constant $M_{\mathcal{C}} < \infty$ such that

$$\|f\|_{m, \infty, \mu^{1/2} \mathbb{1}_{\mathcal{C}}} \leq M_{\mathcal{C}} \|\mu^{1/2} f\|_{m, \infty, \mathbb{1}_{\mathcal{C}}}.$$

for all f such that these norms are defined. If the stronger assumption 3 holds, then this result holds for $\mathcal{C} = \mathcal{D}$ too.

Lemma 6 generalizes lemma A.1 part (d) of Gallant and Nychka (1987) to allow for the weaker assumption 4. Lemma 7 below is analogous to lemma 6, except now using the Sobolev L_2 norm instead of the Sobolev sup-norm. One difference, though, is that the norm on the left hand side now has μ instead of $\mu^{1/2}$.

Lemma 7. Let $\mu : \mathcal{D} \rightarrow \mathbb{R}_+$ be a nonnegative function. Let $m \geq 0$ be an integer. Suppose assumption 4 holds for $\mu = \mu_s$. Then for every compact subset $\mathcal{C} \subseteq \mathcal{D}$, there is a constant $M_{\mathcal{C}} < \infty$ such that

$$\|f\|_{m, 2, \mu \mathbb{1}_{\mathcal{C}}} \leq M_{\mathcal{C}} \|\mu^{1/2} f\|_{m, 2, \mathbb{1}_{\mathcal{C}}}$$

for all f such that these norms are defined. If the stronger assumption 3 holds, then this result holds for $\mathcal{C} = \mathcal{D}$ too.

Lemma 8. Let $\mu : \mathcal{D} \rightarrow \mathbb{R}_+$ be a nonnegative function. Let $m \geq 0$ be an integer. Then there is a constant $M < \infty$ such that

$$\|\mu f\|_{m,\infty} \leq M\|f\|_{m,\infty,\mu}$$

for all functions f such that these norms are defined.

C Some useful lemmas: Proofs

Proof of lemma 1. Let $A \subseteq X$ be $\|\cdot\|_X$ -bounded. Then it is contained in a $\|\cdot\|_X$ -ball. That ball is $\|\cdot\|_Y$ -relatively compact by assumption. So A is a subset of a $\|\cdot\|_Y$ -relatively compact set. Containment is preserved by taking closures of both sets, and hence the $\|\cdot\|_Y$ -closure of A is a subset of a $\|\cdot\|_Y$ -compact set, and is also $\|\cdot\|_Y$ -compact since it is a closed subset of a compact set. \square

Proof of lemma 2. Let $\{a_n\}$ be a sequence in A . Since A is $\|\cdot\|_X$ -closed, any element a such that $\|a_n - a\|_X \rightarrow 0$ must be in A . Let a be such that $\|a_n - a\|_Y \rightarrow 0$. Then $\|a_n - a\|_X \rightarrow 0$ by our norm inequality. Hence $a \in A$. \square

Proof of corollary 1. Follows by repeatedly applying lemma 2. \square

Proof of lemma 3. Let $\{h_n\}$ be a sequence in Ω . Since H_1 is reflexive, and by the Banach-Alaoglu theorem, there is a subsequence $\{h_{n_k}\}$ that weakly converges to some $h \in \Omega$ (e.g., by corollary 1.9.16 of Tao 2010). Let $I : H_1 \rightarrow H_2$ denote the identity operator. Since $H_1 \hookrightarrow H_2$, I is a compact operator. Hence it maps weakly convergent sequences to $\|\cdot\|_2$ -convergent sequences (e.g., proposition 3.3(a) of Conway 1985). That is,

$$\|Ih_{n_k} - Ih\|_2 \rightarrow 0.$$

Hence the image $I\Omega$ is $\|\cdot\|_2$ -sequentially compact. Hence $I\Omega$ is $\|\cdot\|_2$ -closed (e.g., exercise 1.9.11 of Tao 2010). \square

Proof of lemma 4. Since $(X, \|\cdot\|_X)$ is embedded in $(Z, \|\cdot\|_Z)$, there exists a constant $M_1 > 0$ such that

$$\|\cdot\|_Z \leq M_1\|\cdot\|_X.$$

Likewise, by assumption 2, there is a constant constant $M_2 > 0$ such that $\|\cdot\|_Y \leq M_2\|\cdot\|_Z$. Hence

$$\|\cdot\|_Y \leq M_1M_2\|\cdot\|_X.$$

Thus $(X, \|\cdot\|_X)$ is embedded in $(Y, \|\cdot\|_Y)$. Next we need to show that this embedding is compact. Let $A \subseteq X$ be $\|\cdot\|_X$ -bounded. Let $\{a_n\}$ be a sequence in A . By assumption 1 there is a subsequence $\{a_{n_k}\}$ that $\|\cdot\|_Z$ -converges. But by assumption 2, $\|\cdot\|_Z$ is a stronger norm than $\|\cdot\|_Y$ and hence

this subsequence $\|\cdot\|_Y$ -converges. Thus every sequence in A has a $\|\cdot\|_Y$ -convergent subsequence and so A is $\|\cdot\|_Y$ -compact. \square

D Norm inequality lemmas: Proofs

In the proof of lemma 5 and other lemmas, we use the following: The product rule tells us how to differentiate functions like $h(x)g(x)$. The generalization of this rule is called *Leibniz's formula* or the *General Leibniz rule*. For functions u and v that are $|\alpha|$ times continuously differentiable near x , it is

$$[\nabla^\alpha(uv)](x) = \sum_{\{\beta:\beta\leq\alpha\}} \begin{bmatrix} \alpha \\ \beta \end{bmatrix} \nabla^\beta u(x) \nabla^{\alpha-\beta} v(x).$$

Here $\beta \leq \alpha$ is interpreted as being component-wise: $\beta \leq \alpha$ if $\beta_j \leq \alpha_j$ for $1 \leq j \leq d_x$, where d_x is the number of components in the multi-indices β and α , and is also equal to the dimension of the argument x of the functions u and v . Also,

$$\begin{bmatrix} \alpha \\ \beta \end{bmatrix} = \prod_{j=1}^{d_x} \binom{\alpha_j}{\beta_j}$$

where

$$\binom{\alpha_j}{\beta_j} = \frac{\alpha_j!}{\beta_j!(\alpha_j - \beta_j)!}$$

is the binomial coefficient. For a reference on this formula, see Adams and Fournier (2003), page 2.

Proof of lemma 5. Applying Leibniz's formula to the function $\mu(x)^{1/2}f(x)$ we have

$$\nabla^\lambda(\mu^{1/2}f) = \sum_{\{\beta:\beta\leq\lambda\}} \begin{bmatrix} \lambda \\ \beta \end{bmatrix} (\nabla^\beta f)(\nabla^{\lambda-\beta}\mu^{1/2}),$$

for $|\lambda| \leq m + m_0$. By the triangle inequality, this implies

$$\|\nabla^\lambda(\mu^{1/2}f)\|_{0,2,\mathbb{1}_C} \leq \sum_{\{\beta:\beta\leq\lambda\}} \begin{bmatrix} \lambda \\ \beta \end{bmatrix} \|\nabla^{\lambda-\beta}\mu^{1/2}\nabla^\beta f\|_{0,2,\mathbb{1}_C}.$$

Using the bound on the derivatives of $\mu^{1/2}$ we have

$$\begin{aligned}
\|\nabla^{\lambda-\beta}\mu^{1/2}\nabla^\beta f\|_{0,2,\mathbb{1}_C} &= \left(\int_C [\nabla^{\lambda-\beta}\mu^{1/2}(x)\nabla^\beta f(x)]^2 dx \right)^{1/2} \\
&= \left(\int_C |\nabla^{\lambda-\beta}\mu^{1/2}(x)|^2 [\nabla^\beta f(x)]^2 dx \right)^{1/2} \\
&\leq \left(\int_C |K_C\mu^{1/2}(x)|^2 [\nabla^\beta f(x)]^2 dx \right)^{1/2} \\
&= K_C^2 \left(\int_C [\nabla^\beta f(x)]^2 \mu(x) dx \right)^{1/2} \\
&= K_C^2 \|\nabla^\beta f\|_{0,2,\mu\mathbb{1}_C} \\
&\leq K_C^2 \|f\|_{m+m_0,2,\mu\mathbb{1}_C},
\end{aligned}$$

where the last line follows since $m + m_0 \geq 0$. Thus, for $|\lambda| \leq m + m_0$,

$$\|\nabla^\lambda(\mu^{1/2}f)\|_{0,2,\mathbb{1}_C} \leq \left(\sum_{\{\beta:\beta\leq\lambda\}} \begin{bmatrix} \lambda \\ \beta \end{bmatrix} \right) K_C^2 \|f\|_{m+m_0,2,\mu\mathbb{1}_C}.$$

Next,

$$\begin{aligned}
\|\mu^{1/2}f\|_{m+m_0,2,\mathbb{1}_C}^2 &= \sum_{0\leq|\lambda|\leq m+m_0} \|\nabla^\lambda(\mu^{1/2}f)\|_{0,2,\mathbb{1}_C}^2 \\
&\leq \sum_{0\leq|\lambda|\leq m+m_0} \left[\left(\sum_{\{\beta:\beta\leq\lambda\}} \begin{bmatrix} \lambda \\ \beta \end{bmatrix} \right) K_C^2 \|f\|_{m+m_0,2,\mu\mathbb{1}_C} \right]^2 \\
&= \|f\|_{m+m_0,2,\mu\mathbb{1}_C}^2 \left[K_C^2 \sum_{0\leq|\lambda|\leq m+m_0} \left(\sum_{\{\beta:\beta\leq\lambda\}} \begin{bmatrix} \lambda \\ \beta \end{bmatrix} \right) \right]^2 \\
&\equiv \|f\|_{m+m_0,2,\mu\mathbb{1}_C}^2 M_C^2
\end{aligned}$$

and hence

$$\|\mu^{1/2}f\|_{m+m_0,2,\mathbb{1}_C} \leq M_C \|f\|_{m+m_0,2,\mu\mathbb{1}_C}$$

as desired. When assumption 3 holds, the same proof above applies, but the constants now hold over all \mathcal{D} . \square

Proof of lemma 6. We use induction. The inequality holds for $m = 0$ with $M_{\mathcal{C}} = 1$ since

$$\begin{aligned}\|f\|_{0,\infty,\mu^{1/2}\mathbb{1}_{\mathcal{C}}} &= \sup_{x \in \mathcal{D}} |f(x)|\mu^{1/2}(x)\mathbb{1}_{\mathcal{C}}(x) \\ &= \sup_{x \in \mathcal{D}} |\mu^{1/2}(x)f(x)|\mathbb{1}_{\mathcal{C}}(x) \\ &= \|\mu^{1/2}f\|_{0,\infty,\mathbb{1}_{\mathcal{C}}}.\end{aligned}$$

Suppose the inequality holds for m and let $0 < |\lambda| \leq m + 1$. By Leibniz's formula,

$$\nabla^{\lambda}(\mu^{1/2}f) = (\nabla^{\lambda}f)\mu^{1/2} + \sum_{\{\beta:\beta \leq \lambda, \beta \neq \lambda\}} \begin{bmatrix} \lambda \\ \beta \end{bmatrix} (\nabla^{\lambda-\beta}\mu^{1/2})(\nabla^{\beta}f),$$

which implies that

$$\begin{aligned} |(\nabla^{\lambda}f)\mu^{1/2}| &\leq |\nabla^{\lambda}(\mu^{1/2}f)| + \left| \sum_{\{\beta:\beta \leq \lambda, \beta \neq \lambda\}} \begin{bmatrix} \lambda \\ \beta \end{bmatrix} (\nabla^{\lambda-\beta}\mu^{1/2})(\nabla^{\beta}f) \right| \\ &\leq |\nabla^{\lambda}(\mu^{1/2}f)| + K_{\mathcal{C}} \sum_{\{\beta:\beta \leq \lambda, \beta \neq \lambda\}} \begin{bmatrix} \lambda \\ \beta \end{bmatrix} \mu^{1/2}|\nabla^{\beta}f|. \end{aligned}$$

The second line follows by assumption 4, assuming we only evaluate this inequality at $x \in \mathcal{C}$. Taking the supremum over x in \mathcal{C} and the maximum over $|\lambda| \leq m + 1$ gives

$$\|f\|_{m+1,\infty,\mu^{1/2}\mathbb{1}_{\mathcal{C}}} \leq \|\mu^{1/2}f\|_{m+1,\infty,\mathbb{1}_{\mathcal{C}}} + K'_{\mathcal{C}}\|f\|_{m,\infty,\mu^{1/2}\mathbb{1}_{\mathcal{C}}},$$

by the definition of the norms, and since λ isn't included in the sum we get only m derivatives in this last term on the right hand side. Moreover, we picked up an extra \leq since we moved the max and supremum inside the summation in the second term, and then were left with the constant

$$K'_{\mathcal{C}} \equiv K_{\mathcal{C}} \sum_{|\lambda| \leq m} \sum_{\{\beta:\beta \leq \lambda, \beta \neq \lambda\}} \begin{bmatrix} \lambda \\ \beta \end{bmatrix} < \infty.$$

By the induction hypothesis there is an $M'_{\mathcal{C}} < \infty$ such that

$$\|f\|_{m,\infty,\mu^{1/2}\mathbb{1}_{\mathcal{C}}} \leq M'_{\mathcal{C}}\|\mu^{1/2}f\|_{m,\infty,\mathbb{1}_{\mathcal{C}}}.$$

Moreover,

$$\|\mu^{1/2}f\|_{m,\infty,\mathbb{1}_{\mathcal{C}}} \leq \|\mu^{1/2}f\|_{m+1,\infty,\mathbb{1}_{\mathcal{C}}}.$$

Thus

$$\|f\|_{m,\infty,\mu^{1/2}\mathbb{1}_{\mathcal{C}}} \leq M'_{\mathcal{C}}\|\mu^{1/2}f\|_{m+1,\infty,\mathbb{1}_{\mathcal{C}}}.$$

Plugging this into our expression from earlier yields

$$\begin{aligned}
\|f\|_{m+1,\infty,\mu^{1/2}\mathbb{1}_C} &\leq \|\mu^{1/2}f\|_{m+1,\infty,\mathbb{1}_C} + K'_C\|f\|_{m,\infty,\mu^{1/2}\mathbb{1}_C} \\
&\leq \|\mu^{1/2}f\|_{m+1,\infty,\mathbb{1}_C} + K'_CM'_C\|\mu^{1/2}f\|_{m+1,\infty,\mathbb{1}_C} \\
&= (1 + K'_CM'_C)\|\mu^{1/2}f\|_{m+1,\infty,\mathbb{1}_C} \\
&\equiv M_C\|\mu^{1/2}f\|_{m+1,\infty,\mathbb{1}_C}.
\end{aligned}$$

When assumption 3 holds, the same proof above applies, but the constants now hold over all \mathcal{D} . \square

Proof of lemma 7. We will modify the proof of lemma 6 as appropriate. As there, we use proof by induction. For the base case, set $m = 0$. Then

$$\begin{aligned}
\|f\|_{0,2,\mu\mathbb{1}_C} &= \left(\int_C [f(x)]^2 \mu(x) dx \right)^{1/2} \\
&= \left(\int_C [\mu^{1/2}(x)f(x)]^2 dx \right)^{1/2} \\
&= \|\mu^{1/2}f\|_{0,2,\mathbb{1}_C}.
\end{aligned}$$

Thus the result holds for $m = 0$. Now suppose it holds for m . Let $|\lambda|$ be such that $0 < |\lambda| \leq m + 1$. Then, as in the proof of lemma 6, we have

$$\nabla^\lambda(\mu^{1/2}f) = (\nabla^\lambda f)\mu^{1/2} + \sum_{\{\beta:\beta\leq\lambda,\beta\neq\lambda\}} \begin{bmatrix} \lambda \\ \beta \end{bmatrix} (\nabla^{\lambda-\beta}\mu^{1/2})(\nabla^\beta f)$$

by Leibniz's formula. As in that proof, applying our bound on the derivative of the weight function, we get

$$|\nabla^\lambda f|\mu^{1/2} \leq |\nabla^\lambda(\mu^{1/2}f)| + K_C \sum_{\{\beta:\beta\leq\lambda,\beta\neq\lambda\}} \begin{bmatrix} \lambda \\ \beta \end{bmatrix} |\nabla^\beta f|\mu^{1/2}.$$

Now we square both sides and integrate over \mathcal{C} to obtain

$$\begin{aligned}
\int_{\mathcal{C}} |\nabla^\lambda f(x)|^2 \mu(x) dx &\leq \int_{\mathcal{C}} |[\nabla^\lambda(\mu^{1/2} f)](x)|^2 dx \\
&+ \int_{\mathcal{C}} K_{\mathcal{C}}^2 \sum_{\{\tilde{\beta}: \tilde{\beta} \leq \lambda, \tilde{\beta} \neq \lambda\}} \sum_{\{\beta: \beta \leq \lambda, \beta \neq \lambda\}} \begin{bmatrix} \lambda \\ \tilde{\beta} \end{bmatrix} \begin{bmatrix} \lambda \\ \beta \end{bmatrix} |\nabla^{\tilde{\beta}} f(x)| \cdot |\nabla^\beta f(x)| \mu(x) dx \\
&+ \int_{\mathcal{C}} 2|[\nabla^\lambda(\mu^{1/2} f)](x)| K_{\mathcal{C}} \sum_{\{\beta: \beta \leq \lambda, \beta \neq \lambda\}} \begin{bmatrix} \lambda \\ \beta \end{bmatrix} |\nabla^\beta f(x)| \mu^{1/2}(x) dx \\
&= \int_{\mathcal{C}} |[\nabla^\lambda(\mu^{1/2} f)](x)|^2 dx \\
&+ K_{\mathcal{C}}^2 \sum_{\{\tilde{\beta}: \tilde{\beta} \leq \lambda, \tilde{\beta} \neq \lambda\}} \sum_{\{\beta: \beta \leq \lambda, \beta \neq \lambda\}} \begin{bmatrix} \lambda \\ \tilde{\beta} \end{bmatrix} \begin{bmatrix} \lambda \\ \beta \end{bmatrix} \int_{\mathcal{C}} |\nabla^{\tilde{\beta}} f(x)| \cdot |\nabla^\beta f(x)| \mu(x) dx \\
&+ 2K_{\mathcal{C}} \sum_{\{\beta: \beta \leq \lambda, \beta \neq \lambda\}} \begin{bmatrix} \lambda \\ \beta \end{bmatrix} \int_{\mathcal{C}} |[\nabla^\lambda(\mu^{1/2} f)](x)| \cdot |\nabla^\beta f(x)| \mu^{1/2}(x) dx \\
&\equiv (1) + (2) + (3).
\end{aligned}$$

In the third term, we can apply Leibniz's formula again,

$$|\nabla^\beta f| \mu^{1/2} \leq |\nabla^\beta(\mu^{1/2} f)| + K_{\mathcal{C}} \sum_{\{\eta: \eta \leq \beta, \eta \neq \beta\}} \begin{bmatrix} \beta \\ \eta \end{bmatrix} |\nabla^\eta f| \mu^{1/2}$$

to get

$$\begin{aligned}
(3) &\equiv 2K_{\mathcal{C}} \sum_{\{\beta: \beta \leq \lambda, \beta \neq \lambda\}} \begin{bmatrix} \lambda \\ \beta \end{bmatrix} \int_{\mathcal{C}} |[\nabla^\lambda(\mu^{1/2} f)](x)| \cdot |\nabla^\beta f(x)| \mu^{1/2}(x) dx \\
&\leq 2K_{\mathcal{C}} \sum_{\{\beta: \beta \leq \lambda, \beta \neq \lambda\}} \begin{bmatrix} \lambda \\ \beta \end{bmatrix} \left(\int_{\mathcal{C}} |[\nabla^\lambda(\mu^{1/2} f)](x)| \cdot |[\nabla^\beta(\mu^{1/2} f)](x)| dx \right. \\
&\quad \left. + K_{\mathcal{C}} \sum_{\{\eta: \eta \leq \beta, \eta \neq \beta\}} \begin{bmatrix} \beta \\ \eta \end{bmatrix} \int_{\mathcal{C}} |[\nabla^\lambda(\mu^{1/2} f)](x)| \cdot |\nabla^\eta f(x)| \mu^{1/2}(x) dx \right).
\end{aligned}$$

We can apply Leibniz's formula again to eliminate the $|\nabla^\eta f(x)| \mu^{1/2}(x)$ term. Continuing in this manner, we get a sum solely of integrals of the form

$$\int_{\mathcal{C}} |[\nabla^\lambda(\mu^{1/2} f)](x)| \cdot |[\nabla^\beta(\mu^{1/2} f)](x)| dx.$$

Now replace one of the two absolute value terms in the integrand with whichever one is largest.

Suppose its the λ piece. This yields

$$\int_{\mathcal{C}} |[\nabla^\lambda(\mu^{1/2}f)](x)| \cdot |[\nabla^\beta(\mu^{1/2}f)](x)| dx \leq \int_{\mathcal{C}} |[\nabla^\lambda(\mu^{1/2}f)](x)|^2 dx.$$

Thus the third piece is now a sum of terms like this one, where the multi-index in the differential operator can go as high as $|\lambda|$. Summing (3) over $|\lambda|$ with $0 \leq |\lambda| \leq m + 1$ we obtain a sum of many unweighted integrals over \mathcal{C} with integrands of the form $|[\nabla^\lambda(\mu^{1/2}f)](x)|^2$. Now all we have to do is group all these integrals such that our entire expression (3) is a multiple of

$$\sum_{0 \leq |\lambda| \leq m+1} \int_{\mathcal{C}} |[\nabla^\lambda(\mu^{1/2}f)](x)|^2 dx = \|\mu^{1/2}f\|_{m+1,2,\mathbb{1}_{\mathcal{C}}}^2.$$

If there are any ‘missing’ integrals, we can just add on the missing ones (which will give us another inequality, but that’s ok since we only need an upper bound). Thus we see that, after summing over $0 \leq |\lambda| \leq m + 1$, the term (3) is bounded above by

$$C_{3,\mathcal{C}} \|\mu^{1/2}f\|_{m+1,2,\mathbb{1}_{\mathcal{C}}}^2$$

for some constant $C_{3,\mathcal{C}} > 0$.

Consider now the second piece. It is a sum of integrals of the form

$$\int_{\mathcal{C}} |\nabla^{\tilde{\beta}}f(x)| \cdot |\nabla^\beta f(x)|\mu(x) dx.$$

Basically the same argument from third piece applies. We can replace one of the absolute values here with whichever is the largest, thus obtaining an integral of the form

$$\int_{\mathcal{C}} |\nabla^\beta f(x)|^2 \mu(x) dx.$$

Now summing these terms over $0 \leq |\lambda| \leq m + 1$ we see that after grouping all the integrals and adding any missing terms, the entire expression (2) is a multiple of

$$\sum_{0 \leq |\lambda| \leq m} \int_{\mathcal{C}} |\nabla^\lambda f(x)|^2 dx = \|f\|_{m,2,\mathbb{1}_{\mathcal{C}}}^2.$$

It is important here that the sum only goes up to m , not $m + 1$. This is because, in the term (2), the β and $\tilde{\beta}$ pieces are always strictly smaller than λ , and λ itself can only go up to $m + 1$. Hence β and $\tilde{\beta}$ can only go up to m . Thus we see that the term (2) is bounded above by

$$C_{2,\mathcal{C}} \|f\|_{m,2,\mathbb{1}_{\mathcal{C}}}^2$$

for some constant $C_{2,\mathcal{C}} > 0$. Finally, consider the term (1). This term is easy because when we sum

over $0 \leq |\lambda| \leq m + 1$ this term exactly equals

$$\|\mu^{1/2} f\|_{m+1,2,\mathbb{1}_c}^2$$

without having to add any extra terms or mess with the integrands. Combining all these results, we see (by also summing over the left hand side of our original inequality) that

$$\|f\|_{m+1,2,\mu\mathbb{1}_c}^2 \leq (1 + C_{3,c})\|\mu^{1/2} f\|_{m+1,2,\mathbb{1}_c}^2 + C_{2,c}\|f\|_{m,2,\mathbb{1}_c}^2.$$

Now apply the induction hypothesis to the last term to get

$$\begin{aligned} \|f\|_{m+1,2,\mu\mathbb{1}_c}^2 &\leq (1 + C_{3,c})\|\mu^{1/2} f\|_{m+1,2,\mathbb{1}_c}^2 + C_{2,c}\|\mu^{1/2} f\|_{m+1,2,\mathbb{1}_c}^2 \\ &= (1 + C_{3,c} + C_{2,c})\|\mu^{1/2} f\|_{m+1,2,\mathbb{1}_c}^2. \end{aligned}$$

Finally, take the square root of both sides to get

$$\|f\|_{m+1,2,\mu\mathbb{1}_c} \leq (1 + C_{3,c} + C_{2,c})^{1/2}\|\mu^{1/2} f\|_{m+1,2,\mathbb{1}_c}$$

as desired. When assumption 3 holds, the same proof above applies, but the constants now hold over all \mathcal{D} . \square

Proof of lemma 8. As in the proof of lemma 6, we have

$$\nabla^\lambda(\mu^{1/2} f) = (\nabla^\lambda f)\mu^{1/2} + \sum_{\{\beta:\beta \leq \lambda, \beta \neq \lambda\}} \begin{bmatrix} \lambda \\ \beta \end{bmatrix} (\nabla^{\lambda-\beta} \mu^{1/2})(\nabla^\beta f).$$

Hence

$$|\nabla^\lambda(\mu^{1/2} f)| \leq |(\nabla^\lambda f)\mu^{1/2}| + \sum_{\{\beta:\beta \leq \lambda, \beta \neq \lambda\}} \begin{bmatrix} \lambda \\ \beta \end{bmatrix} |(\nabla^\beta f)\mu^{1/2}|.$$

Take the sup over x and the max over $|\lambda| \leq m + 1$ to get

$$\|\mu^{1/2} f\|_{m+1,\infty} \leq \|f\|_{m+1,\infty,\mu^{1/2}} + K'\|f\|_{m,\infty,\mu^{1/2}}.$$

Since $\|f\|_{m,\infty,\mu^{1/2}} \leq \|f\|_{m+1,\infty,\mu^{1/2}}$ we get

$$\|\mu^{1/2} f\|_{m+1,\infty} \leq (1 + K')\|f\|_{m+1,\infty,\mu^{1/2}}.$$

The result follows by evaluating this inequality with the weight μ^2 . \square

E Proof of the compact embedding theorems 1 and 3

In this section we prove theorems 1 and 3. The general outline of the proof of theorem 3 follows the proof of Gallant and Nychka's (1987) lemma A.4, which is a proof of theorem 3 case (1) under the stronger assumption 3.

Proof of theorem 1 (Compact embedding).

1. This follows by the Rellich-Kondrachov theorem (Adams and Fournier (2003) theorem 6.3 part II, equation 5), since m_0 is a positive integer, and since $m_0 > d_x/2$ and \mathcal{D} satisfies the cone condition. In applying the theorem, their j is our m . Their m is our m_0 . Moreover, in their notation, we set $p = 2$ and $k = n = d_x$.
2. This follows by the Rellich-Kondrachov theorem (Adams and Fournier (2003) theorem 6.3, part II, equation 6), since m_0 is a positive integer, and since $m_0 > d_x/2$ and \mathcal{D} satisfies the cone condition. In applying the theorem, as in the previous part above, their j is our m and their m is our m_0 . We set also $q = p = 2$ and $k = n = d_x$.
3. This follows by Adams and Fournier (2003) theorem 1.34 equation 3, and their subsequent remark at the end of that theorem statement.
4. This follows since $\|\cdot\|_{m+m_0,2} \leq M\|\cdot\|_{m+m_0,\infty}$ for some constant $0 < M < \infty$ and hence $\|\cdot\|_{m+m_0,\infty}$ bounded sets are also $\|\cdot\|_{m+m_0,2}$ bounded sets. Then apply part (2), which shows that these bounded sets are $\|\cdot\|_{m,2}$ -relatively compact.
5. This follows by applying the Ascoli-Arzelà theorem; see Adams and Fournier (2003) theorem 1.34 equation 4.

□

Proof of theorem 3 (Compact embedding for unbounded domains with equal weighting). We split the proof into several steps. For each of the cases, define the norms $\|\cdot\|_s$ and $\|\cdot\|_c$ as in table 1.

	$\ \cdot\ _s$	$\ \cdot\ _c$
(1)	$\ \cdot\ _{m+m_0,2,\mu_s}$	$\ \cdot\ _{m,\infty,\mu_c^{1/2}}$
(2)	$\ \cdot\ _{m+m_0,2,\mu_s}$	$\ \cdot\ _{m,2,\mu_c}$
(3)	$\ \cdot\ _{m+m_0,\infty,\mu_s}$	$\ \cdot\ _{m,\infty,\mu_c}$
(4)	$\ \cdot\ _{m+m_0,\infty,\mu_s}$	$\ \cdot\ _{m,2,\mu_c}$

Table 1:

1. **Only look at balls.** By lemma 1, it suffices to show that for any $B > 0$, the $\|\cdot\|_s$ -ball Θ of radius B is $\|\cdot\|_c$ -relatively compact.

$$\text{(Cases 1 and 2.) } \Theta = \{f \in \mathscr{W}_{m+m_0,2,\mu_s}(\mathcal{D}) : \|f\|_{m+m_0,2,\mu_s} \leq B\}.$$

(Cases 3 and 4.) $\Theta = \{f \in \mathcal{C}_{m+m_0, \infty, \mu_s}(\mathcal{D}) : \|f\|_{m+m_0, \infty, \mu_s} \leq B\}$.

2. **Stop worrying about the closure.** We need to show that the $\|\cdot\|_c$ -closure of Θ is $\|\cdot\|_c$ -compact. Let $\{\bar{f}_n\}_{n=1}^\infty$ be a sequence from the $\|\cdot\|_c$ -closure of Θ . It suffices to show that $\{\bar{f}_n\}$ has a convergent subsequence. By the definition of the closure, there exists a sequence $\{f_n\}$ from Θ with

$$\lim_{n \rightarrow \infty} \|f_n - \bar{f}_n\|_c = 0.$$

By the triangle inequality it suffices to show that $\{f_n\}$ has a convergent subsequence. The space

(Case 1.) $\mathcal{C}_{m, \infty, \mu_c^{1/2}}$

(Cases 2 and 4.) $\mathcal{W}_{m, 2, \mu_c}$

(Case 3.) $\mathcal{C}_{m, \infty, \mu_c}$

is complete, so it suffices to show that $\{f_n\}$ has a Cauchy subsequence. The proof of completeness of these spaces is as follows. Recall that a function $f : \mathcal{D} \rightarrow \mathbb{R}$ on the Euclidean domain $\mathcal{D} \subseteq \mathbb{R}^{d_x}$ is locally integrable if for every compact subset $\mathcal{C} \subseteq \mathcal{D}$, $\int_{\mathcal{C}} |f(x)| dx < \infty$. Assumption 6 implies that both $\mu_c^{-1/2}$ (as needed in cases 1, 2, and 4) and μ_c^{-1} (as needed in case 3) are locally integrable on the support of μ_c . Next:

(Case 1) Follows by local integrability of $\mu_c^{-1/2}$ and applying theorem 5.1 of Rodríguez, Álvarez, Romera, and Pestana (2004). To see this, using their notation, assumption 6 ensures that $\Omega_1 = \dots = \Omega_k = \mathbb{R}$ (defined in definition 4 on their page 277) and $\Omega^{(0)} = \mathbb{R}$ (defined on their page 280), and hence by their remark on page 303, the conditions of theorem 5.1 hold. This result is not specific to the one dimensional domain case; for example, see Brown and Opic (1992). The reason we use the power $-1/2$ of μ_c is by the $p = \infty$ case in definition 2 on page 277 of Rodríguez et al. (2004).

(Cases 2 and 4.) Follows by local integrability of $\mu_c^{-1/2}$, and theorem 1.11 of Kufner and Opic (1984) and their remark 4.10 (which extends their theorem to allow for higher order derivatives). The reason we use the power $-1/2$ of μ_c is by the $p = 2 < \infty$ case in definition 2 on page 277 of Rodríguez et al. (2004), or equivalently, equation (1.5) on page 538 of Kufner and Opic (1984).

(Case 3.) Follows by local integrability of μ_c^{-1} and then the same argument as case 1. The reason we use the power -1 of μ_c is by the $p = \infty$ case in definition 2 on page 277 of Rodríguez et al. (2004).

This step is important because functions in the closure may not be differentiable, in which case their norm might not be defined. Even when their norm is defined, functions in the closure do not necessarily satisfy the norm bound. Also, note that if μ_c does not have full support, such as $\mu_c(x) = \mathbb{1}(\|x\|_e \leq M)$ for some constant $M > 0$, then we simply restrict the domain to $\mathcal{D} \cap \{x \in \mathbb{R}^{d_x} : \|x\|_e \leq M\}$ and then proceed as in the bounded support case.

3. **Truncate the domain.** The key idea to deal with the unbounded domain is to partition \mathbb{R}^{d_x} into the open Euclidean ball about the origin

$$\Omega_J = \{x \in \mathbb{R}^{d_x} : x'x < J\} = \{x \in \mathbb{R}^{d_x} : \|x\| < J^2\}$$

and its complement Ω_J^c . As we show in step 9 below, the norm on \mathbb{R}^{d_x} can be split into two pieces: one on Ω_J and another on its complement. We will then show that each of these pieces is small. Restricting ourselves to Ω_J , we will apply existing embedding theorems for bounded domains. We then eventually pick J large enough so that the truncation error is small, which is possible because our weight functions get small as $\|x\|$ gets large.

Let $\mathbb{1}_{\Omega_J}(x) = 1$ if $x \in \Omega_J$ and equal zero otherwise.

4. **Switch to the unweighted norm** so that we can apply an existing compact embedding result for unweighted norms (on bounded domains). Since the f_n are in Θ , we know their weighted norm $\|\cdot\|_s$ is bounded by B . We show that a modified version of the sequence is bounded in an unweighted norm.

(Cases 1 and 2.) The unweighted norm we work with here is $\|\cdot\|_{m+m_0,2,\mathbb{1}_{\Omega_J}}$. For all n ,

$$\begin{aligned} \|\mu_s^{1/2} f_n\|_{m+m_0,2,\mathbb{1}_{\Omega_J}} &\leq M_J \|f_n\|_{m+m_0,2,\mu_s \mathbb{1}_{\Omega_J}} \\ &\leq M_J \|f_n\|_{m+m_0,2,\mu_s} \\ &\leq M_J B. \end{aligned}$$

The first inequality follows by lemma 5, which can be applied by using our assumed bound

$$|\nabla^\lambda \mu_s^{1/2}(x)| \leq K_C \mu_s^{1/2}(x)$$

for all $x \in \mathcal{C}$, where \mathcal{C} is any compact subset of \mathbb{R}^{d_x} . Here and below we let M_J denote the constant from lemma 5 corresponding to the compact set Ω_J . The third inequality follows since $f_n \in \Theta$ and by the definition of Θ . Thus, for each J , $\{\mu_s^{1/2} f_n\}$ is $\|\cdot\|_{m+m_0,2,\mathbb{1}_{\Omega_J}}$ -bounded. Notice that in this step we picked up a power 1/2 of the weight function.

(Case 3.) The unweighted norm we work with here is $\|\cdot\|_{m+m_0,\infty,\mathbb{1}_{\Omega_J}}$. For all n ,

$$\begin{aligned} \|\mu_s f_n\|_{m+m_0,\infty,\mathbb{1}_{\Omega_J}} &\leq M \|f_n\|_{m+m_0,\infty,\mu_s \mathbb{1}_{\Omega_J}} \\ &\leq M \|f_n\|_{m+m_0,\infty,\mu_s} \\ &\leq MB. \end{aligned}$$

The first inequality follows by lemma 8. The third inequality follows since $f_n \in \Theta$ and by the definition of Θ . Thus, for each J , $\{\mu_s f_n\}$ is $\|\cdot\|_{m+m_0,\infty,\mathbb{1}_{\Omega_J}}$ -bounded.

(Case 4.) The unweighted norm we work with here is $\|\cdot\|_{m+m_0,\infty,\mathbb{1}_{\Omega_J}}$. For all n ,

$$\begin{aligned}
\|\mu_s^{1/2} f_n\|_{m+m_0,\infty,\mathbb{1}_{\Omega_J}} &\leq M \|f_n\|_{m+m_0,\infty,\mu_s^{1/2}\mathbb{1}_{\Omega_J}} \\
&= M \max_{0\leq|\lambda|\leq m+m_0} \sup_{x\in\mathcal{D}} |\nabla^\lambda f_n(x)| \mu_s^{1/2}(x) \mathbb{1}_{\Omega_J}(x) \\
&= M \max_{0\leq|\lambda|\leq m+m_0} \sup_{x\in\mathcal{D}} |\nabla^\lambda f_n(x)| \mu_s(x) \mu_s^{-1/2}(x) \mathbb{1}_{\Omega_J}(x) \\
&\leq M \left(\max_{0\leq|\lambda|\leq m+m_0} \sup_{x\in\mathcal{D}} |\nabla^\lambda f_n(x)| \mu_s(x) \right) \sup_{\|x\|_e > J^2} \mu_s^{-1/2}(x) \\
&= M \|f_n\|_{m+m_0,\infty,\mu_s^{1/2}} \sup_{\|x\|_e > J^2} \mu_s^{-1/2}(x) \\
&\leq MBK_J.
\end{aligned}$$

The first inequality follows by lemma 8. The final inequality follows since $f_n \in \Theta$ and by the definition of Θ , as well as by assumption that μ_s is bounded away from zero for any compact subset of \mathbb{R}^{d_x} . Thus, for each J , $\{\mu_s^{1/2} f_n\}$ is $\|\cdot\|_{m+m_0,\infty,\mathbb{1}_{\Omega_J}}$ -bounded.

5. Apply an embedding theorem for bounded domains.

(Case 1.) By theorem 1 part 1, $\mathscr{W}_{m+m_0,2,\mathbb{1}_{\Omega_J}}$ is compactly embedded in $\mathcal{C}_{m,\infty,\mathbb{1}_{\Omega_J}}$. Thus, since $\{\mu_s^{1/2} f_n\}$ is $\|\cdot\|_{m+m_0,2,\mathbb{1}_{\Omega_J}}$ -bounded, it is relatively compact in $\mathcal{C}_{m,\infty,\mathbb{1}_{\Omega_J}}$.

(Case 2.) By theorem 1 part 2, $\mathscr{W}_{m+m_0,2,\mathbb{1}_{\Omega_J}}$ is compactly embedded in $\mathscr{W}_{m,2,\mathbb{1}_{\Omega_J}}$. Thus, since $\{\mu_s^{1/2} f_n\}$ is $\|\cdot\|_{m+m_0,2,\mathbb{1}_{\Omega_J}}$ -bounded, it is relatively compact in $\mathscr{W}_{m,2,\mathbb{1}_{\Omega_J}}$.

(Case 3.) By theorem 1 part 3, $\mathcal{C}_{m+m_0,\infty,\mathbb{1}_{\Omega_J}}$ is compactly embedded in $\mathcal{C}_{m,\infty,\mathbb{1}_{\Omega_J}}$. Thus, since $\{\mu_s f_n\}$ is $\|\cdot\|_{m+m_0,\infty,\mathbb{1}_{\Omega_J}}$ -bounded, it is relatively compact in $\mathcal{C}_{m,\infty,\mathbb{1}_{\Omega_J}}$.

(Case 4.) By theorem 1 part 4, $\mathcal{C}_{m+m_0,\infty,\mathbb{1}_{\Omega_J}}$ is compactly embedded in $\mathscr{W}_{m,2,\mathbb{1}_{\Omega_J}}$. Thus, since $\{\mu_s^{1/2} f_n\}$ is $\|\cdot\|_{m+m_0,\infty,\mathbb{1}_{\Omega_J}}$ -bounded, it is relatively compact in $\mathscr{W}_{m,2,\mathbb{1}_{\Omega_J}}$.

In cases 1, 2, and 4 we used that $m_0 > d_x/2$, and note that Ω_J satisfies the cone condition. In case 3 we used that Ω_J is convex and $m_0 \geq 1$.

6. Extract a subsequence. Set $J = 1$. By the previous step, there is a subsequence

(Case 1.) $\{\mu_s^{1/2} f_j^{(1)}\}_{j=1}^\infty$ and a ψ_1 in $\mathcal{C}_{m,\infty,\mathbb{1}_{\Omega_1}}$ such that

$$\lim_{j \rightarrow \infty} \|\mu_s^{1/2} f_j^{(1)} - \psi_1\|_{m,\infty,\mathbb{1}_{\Omega_1}} = 0.$$

(Cases 2 and 4.) $\{\mu_s^{1/2} f_j^{(1)}\}_{j=1}^\infty$ and a ψ_1 in $\mathcal{W}_{m,2,1\Omega_1}$ such that

$$\lim_{j \rightarrow \infty} \|\mu_s^{1/2} f_j^{(1)} - \psi_1\|_{m,2,1\Omega_1} = 0.$$

(Case 3.) $\{\mu_s f_j^{(1)}\}_{j=1}^\infty$ and a ψ_1 in $\mathcal{C}_{m,\infty,1\Omega_1}$ such that

$$\lim_{j \rightarrow \infty} \|\mu_s f_j^{(1)} - \psi_1\|_{m,\infty,1\Omega_1} = 0.$$

7. **Do it for all J .** Repeating this argument for all J , we have a bunch of nested subsequences

(Cases 1, 2, and 4.)

$$\{\mu_s^{1/2} f_n\} \supset \{\mu_s^{1/2} f_j^{(1)}\} \supset \{\mu_s^{1/2} f_j^{(2)}\} \supset \dots$$

each with

(Case 1.)

$$\lim_{j \rightarrow \infty} \|\mu_s^{1/2} f_j^{(J)} - \psi_J\|_{m,\infty,1\Omega_J} = 0.$$

(Cases 2 and 4.)

$$\lim_{j \rightarrow \infty} \|\mu_s^{1/2} f_j^{(J)} - \psi_J\|_{m,2,1\Omega_J} = 0.$$

(Case 3.)

$$\{\mu_s f_n\} \supset \{\mu_s f_j^{(1)}\} \supset \{\mu_s f_j^{(2)}\} \supset \dots$$

each with

$$\lim_{j \rightarrow \infty} \|\mu_s f_j^{(J)} - \psi_J\|_{m,\infty,1\Omega_J} = 0.$$

The reason we have to extract a further subsequence from

(Cases 1, 2, and 4.) $\{\mu_s^{1/2} f_1^{(1)}\}$ is that $\{\mu_s^{1/2} f_1^{(1)}\}$

(Case 3.) $\{\mu_s f_1^{(1)}\}$ is that $\{\mu_s f_1^{(1)}\}$

only converges in the norm with $J = 1$; it may not converge in the norm with $J = 2$. So we extract a further subsequence which does converge in the norm with $J = 2$, and so on.

8. **Define the main subsequence.** Set $f_j = f_j^{(j)}$. Then $\{f_j\}$ is a subsequence of $\{f_n\}$. Our goal is to show that $\{f_j\}$ is $\|\cdot\|_c$ -Cauchy. Let $\varepsilon > 0$ be given. This is a kind of diagonalization argument.

9. **Split the consistency norm into two pieces.**

(Cases 1 and 3.) For any weight μ_c and any set Ω , we have

$$\begin{aligned}
\|f\|_{m,\infty,\mu_c} &\equiv \max_{0 \leq |\lambda| \leq m} \sup_{x \in \mathbb{R}^{d_x}} |\nabla^\lambda f(x)| \mu_c(x) (\mathbb{1}_\Omega(x) + \mathbb{1}_{\Omega^c}(x)) \\
&= \max_{0 \leq |\lambda| \leq m} \sup_{x \in \mathbb{R}^{d_x}} \left(|\nabla^\lambda f(x)| \mu_c(x) \mathbb{1}_\Omega(x) + |\nabla^\lambda f(x)| \mu_c(x) \mathbb{1}_{\Omega^c}(x) \right) \\
&\leq \max_{0 \leq |\lambda| \leq m} \sup_{x \in \mathbb{R}^{d_x}} |\nabla^\lambda f(x)| \mu_c(x) \mathbb{1}_\Omega(x) + \max_{0 \leq |\lambda| \leq m} \sup_{x \in \mathbb{R}^{d_x}} |\nabla^\lambda f(x)| \mu_c(x) \mathbb{1}_{\Omega^c}(x) \\
&= \|f\|_{m,\infty,\mu_c \mathbb{1}_\Omega} + \|f\|_{m,\infty,\mu_c \mathbb{1}_{\Omega^c}},
\end{aligned}$$

where Ω^c is the complement of Ω . Hence, for any J , and for any f_j and f_k in our main subsequence $\{f_j\}$ we have

(Case 1.)

$$\|f_j - f_k\|_{m,\infty,\mu_c^{1/2}} \leq \|f_j - f_k\|_{m,\infty,\mu_c^{1/2} \mathbb{1}_{\Omega_J}} + \|f_j - f_k\|_{m,\infty,\mu_c^{1/2} \mathbb{1}_{\Omega_J^c}}.$$

(Case 3.)

$$\|f_j - f_k\|_{m,\infty,\mu_c} \leq \|f_j - f_k\|_{m,\infty,\mu_c \mathbb{1}_{\Omega_J}} + \|f_j - f_k\|_{m,\infty,\mu_c \mathbb{1}_{\Omega_J^c}}.$$

(Cases 2 and 4.) We want to show that

$$\|f\|_{m,2,\mu_c} \leq \|f\|_{m,2,\mu_c \mathbb{1}_{\Omega_J}} + \|f\|_{m,2,\mu_c \mathbb{1}_{\Omega_J^c}}.$$

We have

$$\begin{aligned}
\|f\|_{m,2,\mu_c}^2 &= \sum_{0 \leq |\lambda| \leq m} \int [\nabla^\lambda f(x)]^2 \mu_c(x) dx \\
&= \sum_{0 \leq |\lambda| \leq m} \left[\int [\nabla^\lambda f(x)]^2 \mu_c(x) \mathbb{1}_{\Omega_J}(x) dx + \int [\nabla^\lambda f(x)]^2 \mu_c(x) \mathbb{1}_{\Omega_J^c}(x) dx \right] \\
&= \sum_{0 \leq |\lambda| \leq m} \int [\nabla^\lambda f(x)]^2 \mu_c(x) \mathbb{1}_{\Omega_J}(x) dx + \sum_{0 \leq |\lambda| \leq m} \int [\nabla^\lambda f(x)]^2 \mu_c(x) \mathbb{1}_{\Omega_J^c}(x) dx \\
&= \|f\|_{m,2,\mu_c \mathbb{1}_{\Omega_J}}^2 + \|f\|_{m,2,\mu_c \mathbb{1}_{\Omega_J^c}}^2.
\end{aligned}$$

Hence

$$\begin{aligned}
\|f\|_{m,2,\mu_c} &= \sqrt{\|f\|_{m,2,\mu_c \mathbb{1}_{\Omega_J}}^2 + \|f\|_{m,2,\mu_c \mathbb{1}_{\Omega_J^c}}^2} \\
&\leq \|f\|_{m,2,\mu_c \mathbb{1}_{\Omega_J}} + \|f\|_{m,2,\mu_c \mathbb{1}_{\Omega_J^c}},
\end{aligned}$$

where the last line follows by $\sqrt{a^2 + b^2} \leq a + b$ for $a, b \geq 0$. Hence, for any J , and for

any f_j and f_k in our main subsequence $\{f_j\}$ we have

$$\|f_j - f_k\|_{m,2,\mu_c} \leq \|f_j - f_k\|_{m,2,\mu_c \mathbb{1}_{\Omega_J}} + \|f_j - f_k\|_{m,2,\mu_c \mathbb{1}_{\Omega_J^c}},$$

where recall that Ω_J^c is the complement of Ω_J .

Now we just need to show that if j, k are sufficiently far out in the sequence, and J is large enough, that both of these pieces on the right hand side are small.

10. Outside truncation piece is small.

(Case 1.) Since $f_j \in \Theta$ for all j , $\|f_j\|_{m+m_0,2,\mu_s} \leq B$ for all j . This combined with assumption 5 let us apply lemma 9 to find a large enough J such that

$$\|f_j\|_{m,\infty,\mu_c^{1/2} \mathbb{1}_{\Omega_J^c}} < \frac{\varepsilon}{4}$$

for all j . By the triangle inequality,

$$\|f_j - f_k\|_{m,\infty,\mu_c^{1/2} \mathbb{1}_{\Omega_J^c}} < 2\frac{\varepsilon}{4} = \frac{\varepsilon}{2}.$$

(Case 2.) For this case,

$$\begin{aligned} \|f_j\|_{m,2,\mu_s \mathbb{1}_{\Omega_J^c}} &\leq \|f_j\|_{m,2,\mu_s} \\ &\leq \|f_j\|_{m+m_0,2,\mu_s} \\ &\leq B, \end{aligned}$$

where the last line follows since $f_j \in \Theta$. Next, by assumption 1,

$$\frac{\mu_c(x)}{\mu_s(x)} \rightarrow 0 \quad \text{as} \quad x'x \rightarrow \infty.$$

So we can choose J large enough that

$$\left(\frac{\mu_c(x)}{\mu_s(x)}\right)^{1/2} < \frac{\varepsilon^2}{4^2 B^2}$$

for all $x'x > J$; i.e., for all $x \in \Omega_J^c$. Next, we have

$$\begin{aligned}
\|f_j\|_{m,2,\mu_c\mathbb{1}_{\Omega_J^c}}^2 &= \sum_{0 \leq |\lambda| \leq m} \int_{\Omega_J^c} |\nabla^\lambda f_j(x)|^2 \mu_c(x) dx \\
&= \sum_{0 \leq |\lambda| \leq m} \int_{\Omega_J^c} |\nabla^\lambda f_j(x)|^2 \mu_s(x) \frac{\mu_c(x)}{\mu_s(x)} dx \\
&\leq \sum_{0 \leq |\lambda| \leq m} \int_{\Omega_J^c} |\nabla^\lambda f_j(x)|^2 \mu_s(x) \frac{\varepsilon^2}{4^2 M} dx \\
&= \frac{\varepsilon^2}{4^2 B^2} \sum_{0 \leq |\lambda| \leq m} \int_{\Omega_J^c} |\nabla^\lambda f_j(x)|^2 \mu_s(x) dx \\
&= \frac{\varepsilon^2}{4^2 B^2} \|f_j\|_{m,2,\mu_s\mathbb{1}_{\Omega_J^c}}^2 \\
&\leq \frac{\varepsilon}{4^2 B^2} B^2 \\
&= \frac{\varepsilon^2}{4^2}.
\end{aligned}$$

(Case 4.) For this case,

$$\begin{aligned}
\|f_j\|_{m,2,\mu_c\mathbb{1}_{\Omega_J^c}}^2 &= \sum_{0 \leq |\lambda| \leq m} \int_{\Omega_J^c} |\nabla^\lambda f_j(x)|^2 \mu_c(x) dx \\
&= \sum_{0 \leq |\lambda| \leq m} \int_{\Omega_J^c} |\nabla^\lambda f_j(x)|^2 \mu_s^2(x) \frac{\mu_c(x)}{\mu_s^2(x)} dx \\
&\leq C \|f\|_{m,\infty,\mu_s}^2 \int_{\Omega_J^c} \frac{\mu_c(x)}{\mu_s^2(x)} dx \\
&\leq C B^2 \int_{\Omega_J^c} \frac{\mu_c(x)}{\mu_s^2(x)} dx \\
&\leq \frac{\varepsilon^2}{4^2},
\end{aligned}$$

where in the last step we choose J large enough so that¹

$$\int_{\Omega_J^c} \frac{\mu_c(x)}{\mu_s^2(x)} dx \leq \frac{\varepsilon^2}{4^2 C B^2}.$$

This is possible by our assumption that the integral on the left hand side is finite for at least some J . That implies, by the monotone convergence theorem for sequences of pointwise decreasing functions (e.g., Folland (1999) exercise 15 on page 52), that the integral converges to zero as $J \rightarrow \infty$.

¹Here we see that we could weaken our assumption on the integral to merely that $\int_{\Omega_J^c} \mu_c(x)/\mu_s(x) dx < \infty$ for some J if we switched to using the weight $\mu_s^{1/2}$ instead of μ_s in defining the parameter space.

(Cases 2 and 4). Take the square root of both sides to get

$$\|f_j\|_{m,2,\mu_c \mathbb{1}_{\Omega_J^c}} \leq \frac{\varepsilon}{4}.$$

By the triangle inequality,

$$\|f_j - f_k\|_{m,2,\mu_c \mathbb{1}_{\Omega_J^c}} < 2\frac{\varepsilon}{4} = \frac{\varepsilon}{2}.$$

(Case 3.) We have

$$\begin{aligned} \|f_j\|_{m,\infty,\mu_c \mathbb{1}_{\Omega_J^c}} &= \max_{0 \leq |\lambda| \leq m} \sup_{x \in \mathcal{D}} |\nabla^\lambda f_j(x)| \mu_c(x) \mathbb{1}_{\Omega_J^c}(x) \\ &= \max_{0 \leq |\lambda| \leq m} \sup_{x \in \mathcal{D}} |\nabla^\lambda f_j(x)| \mu_s(x) \frac{\mu_c(x)}{\mu_s(x)} \mathbb{1}_{\Omega_J^c}(x) \\ &\leq \left(\max_{0 \leq |\lambda| \leq m} \sup_{x \in \mathcal{D}} |\nabla^\lambda f_j(x)| \mu_s(x) \right) \sup_{\|x\|_e \geq J^2} \frac{\mu_c(x)}{\mu_s(x)} \\ &= \|f_j\|_{m,\infty,\mu_s} \sup_{\|x\|_e \geq J^2} \frac{\mu_c(x)}{\mu_s(x)} \\ &\leq \|f_j\|_{m+m_0,\infty,\mu_s} \sup_{\|x\|_e \geq J^2} \frac{\mu_c(x)}{\mu_s(x)} \\ &\leq B \sup_{\|x\|_e \geq J^2} \frac{\mu_c(x)}{\mu_s(x)} \\ &\leq B \frac{\varepsilon}{4B} \\ &= \frac{\varepsilon}{4}. \end{aligned}$$

The second to last line follows by choosing J large enough, and using assumption 1. By the triangle inequality,

$$\|f_j - f_k\|_{m,\infty,\mu_c \mathbb{1}_{\Omega_J^c}} < 2\frac{\varepsilon}{4} = \frac{\varepsilon}{2}.$$

11. **Inside truncation piece is small.** In the previous step we chose a specific value of J , so here we take J as fixed. $\{f_j\}_{j=J}^\infty = \{f_j^{(j)}\}_{j=J}^\infty$ (equality follows by definition of f_j) is a subsequence from $\{f_j^{(J)}\}$. This follows since the subsequences are nested:

$$\text{(Cases 1, 2, and 4.) } \{\mu_s^{1/2} f_n\} \supset \{\mu_s^{1/2} f_j^{(1)}\} \supset \{\mu_s^{1/2} f_j^{(2)}\} \supset \dots$$

$$\text{(Case 3.) } \{\mu_s f_n\} \supset \{\mu_s f_j^{(1)}\} \supset \{\mu_s f_j^{(2)}\} \supset \dots$$

(Case 1.) Since $\{\mu_s^{1/2} f_j^{(J)}\}$ converges in the norm $\|\cdot\|_{m,\infty,\mathbb{1}_{\Omega_J}}$ it is also Cauchy in that norm. Thus there is some K large enough (take $K > J$) such that

$$\|\mu_s^{1/2}(f_j - f_k)\|_{m,\infty,\mathbb{1}_{\Omega_J}} < \frac{\varepsilon}{2M_5^{1/2}M'_J}$$

for all $k, j > K$. Here M'_J is the constant from applying lemma 6 to $\mathcal{C} = \Omega_J$. Notice that this constant is different from M_J , which comes from applying lemma 5.

Hence

$$\begin{aligned}
\|f_j - f_k\|_{m, \infty, \mu_c^{1/2} \mathbb{1}_{\Omega_J}} &\leq M_5^{1/2} \|f_j - f_k\|_{m, \infty, \mu_s^{1/2} \mathbb{1}_{\Omega_J}} \\
&\leq M_5^{1/2} M'_J \|\mu_s^{1/2}(f_j - f_k)\|_{m, \infty, \mathbb{1}_{\Omega_J}} && \text{by lemma 6} \\
&< M_5^{1/2} M'_J \frac{\varepsilon}{2M_5^{1/2} M'_J} \\
&= \frac{\varepsilon}{2}.
\end{aligned}$$

Applying lemma 6 uses assumption 4. The first line follows since

$$\begin{aligned}
\|f\|_{m, \infty, \mu_c^{1/2} \mathbb{1}_{\Omega_J}} &= \max_{0 \leq |\lambda| \leq m} \sup_{x \in \mathbb{R}^{d_x}} |\nabla^\lambda f(x)| \mu_c^{1/2}(x) \mathbb{1}_{\Omega_J}(x) \\
&= \max_{0 \leq |\lambda| \leq m} \sup_{x \in \mathbb{R}^{d_x}} |\nabla^\lambda f(x)| \mu_s^{1/2}(x) \left(\frac{\mu_c(x)}{\mu_s(x)} \right)^{1/2} \mathbb{1}_{\Omega_J}(x) \\
&\leq \max_{0 \leq |\lambda| \leq m} \sup_{x \in \mathbb{R}^{d_x}} |\nabla^\lambda f(x)| \mu_s^{1/2}(x) M_5^{1/2} \mathbb{1}_{\Omega_J}(x) \\
&= M_5^{1/2} \|f\|_{m, \infty, \mu_s^{1/2} \mathbb{1}_{\Omega_J}},
\end{aligned}$$

where we used our assumption 2 that

$$\frac{\mu_c(x)}{\mu_s(x)} \leq M_5$$

for all $x \in \mathbb{R}^{d_x}$.

(Cases 2 and 4.) Since $\{\mu_s^{1/2} f_j^{(J)}\}$ converges in the norm $\|\cdot\|_{m, 2, \mathbb{1}_{\Omega_J}}$ it is also Cauchy in that norm. Thus there is a K large enough (take $K > J$) such that

$$\|\mu_s^{1/2}(f_j - f_k)\|_{m, 2, \mathbb{1}_{\Omega_J}} < \frac{\varepsilon}{2M_5^{1/2} M'_J}$$

for all $j, k > K$. Here M'_J is the constant from applying lemma 7 to $\mathcal{C} = \Omega_J$. Applying this lemma uses assumption 4. We need to show that this implies

$$\|f_j - f_k\|_{m, 2, \mu_c \mathbb{1}_{\Omega_J}}$$

is small ($\leq \varepsilon/2$) for all $j, k > K$. We have

$$\begin{aligned}
\|f\|_{m,2,\mu_c\mathbb{1}_{\Omega_J}} &= \left(\sum_{0 \leq |\lambda| \leq m} \int_{\mathcal{D}} [\nabla^\lambda f(x)]^2 \mu_c(x) \mathbb{1}_{\Omega_J}(x) dx \right)^{1/2} \\
&= \left(\sum_{0 \leq |\lambda| \leq m} \int_{\mathcal{D}} [\nabla^\lambda f(x)]^2 \mu_s(x) \frac{\mu_c(x)}{\mu_s(x)} \mathbb{1}_{\Omega_J}(x) dx \right)^{1/2} \\
&\leq \left(\sup_{x \in \mathbb{R}^{d_x}} \frac{\mu_c(x)}{\mu_s(x)} \sum_{0 \leq |\lambda| \leq m} \int_{\mathcal{D}} [\nabla^\lambda f(x)]^2 \mu_s(x) \mathbb{1}_{\Omega_J}(x) dx \right)^{1/2} \\
&\leq M_5^{1/2} \left(\sum_{0 \leq |\lambda| \leq m} \int_{\mathcal{D}} [\nabla^\lambda f(x)]^2 \mu_s(x) \mathbb{1}_{\Omega_J}(x) dx \right)^{1/2} \\
&= M_5^{1/2} \|f\|_{m,2,\mu_s\mathbb{1}_{\Omega_J}},
\end{aligned}$$

where the fourth line follows by assumption 2, which said that

$$\frac{\mu_c(x)}{\mu_s(x)} \leq M_5$$

for all $x \in \mathbb{R}^{d_x}$. This shows us how to switch from weighting with μ_c to weighting with μ_s . By lemma 7,

$$\|f\|_{m,2,\mu_s\mathbb{1}_{\Omega_J}} \leq M'_J \|\mu_s^{1/2} f\|_{m,2,\mathbb{1}_{\Omega_J}}.$$

Thus we are done since

$$\begin{aligned}
\|f_j - f_k\|_{m,2,\mu_c\mathbb{1}_{\Omega_J}} &\leq M_5^{1/2} \|f_j - f_k\|_{m,2,\mu_s\mathbb{1}_{\Omega_J}} \\
&\leq M_5^{1/2} M'_J \|\mu_s^{1/2} (f_j - f_k)\|_{m,2,\mathbb{1}_{\Omega_J}} \\
&\leq M_5^{1/2} M'_J \frac{\varepsilon}{2M_5^{1/2} M'_J} \\
&= \frac{\varepsilon}{2}.
\end{aligned}$$

(Case 3.) Since $\{\mu_s f_j^{(J)}\}$ converges in the norm $\|\cdot\|_{m,\infty,\mathbb{1}_{\Omega_J}}$ it is also Cauchy in that norm. Thus there is some K large enough (take $K > J$) such that

$$\|\mu_s(f_j - f_k)\|_{m,\infty,\mathbb{1}_{\Omega_J}} < \frac{\varepsilon}{2M_5 M'_J}$$

for all $k, j > K$. Here M'_J is the constant from applying lemma 6 to $\mathcal{C} = \Omega_J$. Notice that this constant is different from M_J , which comes from applying lemma 5.

Hence

$$\begin{aligned}
\|f_j - f_k\|_{m,\infty,\mu_c\mathbb{1}_{\Omega_J}} &\leq M_5\|f_j - f_k\|_{m,\infty,\mu_s\mathbb{1}_{\Omega_J}} \\
&\leq M_5M'_J\|\mu_s(f_j - f_k)\|_{m,\infty,\mathbb{1}_{\Omega_J}} \quad \text{by lemma 6 applied with } \mu = \mu_s^2 \\
&< M_5M'_J\frac{\varepsilon}{2M_5M'_J} \\
&= \frac{\varepsilon}{2}.
\end{aligned}$$

Applying lemma 6 uses assumption 4. The first line follows since

$$\begin{aligned}
\|f\|_{m,\infty,\mu_c\mathbb{1}_{\Omega_J}} &= \max_{0\leq|\lambda|\leq m} \sup_{x\in\mathcal{D}} |\nabla^\lambda f(x)|\mu_c(x)\mathbb{1}_{\Omega_J}(x) \\
&= \max_{0\leq|\lambda|\leq m} \sup_{x\in\mathcal{D}} |\nabla^\lambda f(x)|\mu_s(x)\frac{\mu_c(x)}{\mu_s(x)}\mathbb{1}_{\Omega_J}(x) \\
&\leq \max_{0\leq|\lambda|\leq m} \sup_{x\in\mathcal{D}} |\nabla^\lambda f(x)|\mu_s(x)M_5\mathbb{1}_{\Omega_J}(x) \\
&= M_5\|f\|_{m,\infty,\mu_s\mathbb{1}_{\Omega_J}},
\end{aligned}$$

where the third line follows by assumption 2.

12. **Put previous two steps together.** We now have

$$\|f_j - f_k\|_c \leq \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon$$

for all $k, j > K$. The constants only depend on the choice of weight functions, not J or any other variable that changes along the sequence. Thus we have shown that $\{f_j\}$ is $\|\cdot\|_c$ -Cauchy. □

Lemma 9. Let $\mu_c, \mu_s : \mathcal{D} \rightarrow \mathbb{R}_+$ be nonnegative functions. Let $m, m_0 \geq 0$ be integers. Let Ω_J be defined as in the proof of either theorem 3 or 5. Suppose assumption 5 holds and $\|f\|_{m+m_0,2,\mu_s} \leq B$. Then there is a function $K(J)$ such that

$$\|f\|_{m,\infty,\mu_c^{1/2}\mathbb{1}_{\Omega_J^c}} \leq K(J)$$

where $K(J) \rightarrow 0$ as $J \rightarrow \infty$.

Proof of lemma 9. For all $0 \leq |\lambda| \leq m$,

$$\begin{aligned}
\|\nabla^\lambda f\|_{0,\infty,\mu_c^{1/2}\mathbb{1}_{\Omega_j^c}} &= \sup_{x \in \Omega_j^c} |\nabla^\lambda f(x)| \mu_c^{1/2}(x) \\
&= \sup_{x \in \Omega_j^c} |\nabla^\lambda f(x)| \tilde{\mu}_c^{1/2}(x) \frac{1}{g(x)} \\
&\leq \sup_{x \in \Omega_j^c} |\nabla^\lambda f(x)| \tilde{\mu}_c^{1/2}(x) \sup_{x \in \Omega_j^c} \frac{1}{g(x)} \\
&= \|\tilde{\mu}_c^{1/2} \nabla^\lambda f\|_{0,\infty,\mathbb{1}_{\Omega_j^c}} \sup_{x \in \Omega_j^c} \frac{1}{g(x)} \\
&\leq \|\tilde{\mu}_c^{1/2} \nabla^\lambda f\|_{0,\infty} \sup_{x \in \Omega_j^c} \frac{1}{g(x)}.
\end{aligned}$$

By the Sobolev embedding theorem (Adams and Fournier 2003, theorem 4.12, part 1, case A, equation 1) there is a constant $M_2 < \infty$ such that

$$\|g\|_{0,\infty} \leq M_2 \|g\|_{m_0,2}$$

for all g in $\mathcal{W}_{m_0,2}$ where $m_0 > d_x/2$. This inequality implies

$$\begin{aligned}
\|\tilde{\mu}_c^{1/2} \nabla^\lambda f\|_{0,\infty} &\leq M_2 \|\tilde{\mu}_c^{1/2} \nabla^\lambda f\|_{m_0,2} \\
&\leq M_2 M \|\nabla^\lambda f\|_{m_0,2,\mu_s} \\
&\equiv M_3 \|\nabla^\lambda f\|_{m_0,2,\mu_s}.
\end{aligned}$$

The second line follows by using assumption 5 in arguments as in the proof of lemma 5. Hence

$$\begin{aligned}
\|\nabla^\lambda f\|_{0,\infty,\mu_c^{1/2}\mathbb{1}_{\Omega_j^c}} &\leq M_3 \|\nabla^\lambda f\|_{m_0,2,\mu_s} \sup_{x \in \Omega_j^c} \frac{1}{g(x)} \\
&\leq M_3 \left(\sum_{0 \leq |\eta| \leq |\lambda| + m_0} \|\nabla^\eta f\|_{0,2,\mu_s} \right) \sup_{x \in \Omega_j^c} \frac{1}{g(x)} \\
&\leq M_3 \left(\sum_{0 \leq |\eta| \leq |\lambda| + m_0} \|f\|_{m+m_0,2,\mu_s} \right) \sup_{x \in \Omega_j^c} \frac{1}{g(x)} \\
&\leq M_3 \left(\sum_{0 \leq |\eta| \leq |\lambda| + m_0} B \right) \sup_{x \in \Omega_j^c} \frac{1}{g(x)} \\
&\leq M_3 \left(\sum_{0 \leq |\eta| \leq m+m_0} B \right) \sup_{x \in \Omega_j^c} \frac{1}{g(x)} \\
&\equiv K(J).
\end{aligned}$$

The second line uses $\sqrt{a_1^2 + \dots + a_n^2} \leq a_1 + \dots + a_n$ and the definition of the Sobolev L_2 norm. The

third line uses $|a_i| \leq \sqrt{a_1^2 + \dots + a_n^2}$ for $i = 1, \dots, n$. By the definition of Ω_J , and since $g(x) \rightarrow \infty$ as $\|x\|_e \rightarrow \infty$ (for $\mathcal{D} = \mathbb{R}^{d_x}$) or as x approaches $\text{Bd}(\overline{\mathcal{D}})$ (for bounded \mathcal{D}),

$$\sup_{x \in \Omega_J^c} \frac{1}{g(x)} \rightarrow 0.$$

Hence $K(J) \rightarrow 0$ as $J \rightarrow \infty$. Finally,

$$\begin{aligned} \|f\|_{m, \infty, \mu_c^{1/2} \mathbb{1}_{\Omega_J^c}} &= \max_{0 \leq |\lambda| \leq m} \|\nabla^\lambda f\|_{0, \infty, \mu_c^{1/2} \mathbb{1}_{\Omega_J^c}} \\ &\leq K(J). \end{aligned}$$

□

F Proofs of the compact embedding theorems 5 and 7

Proof of theorem 5 (Compact embedding for unbounded domains with product weighting). For cases 1–3, we apply lemma S1 below, which allows us to convert our previous compact embedding and closedness results for equal weighting to results for product weighting. For case 4, we do not have such a prior result because it's not clear how to define equal weighted Hölder norms, as discussed in the main paper. Hence for this case we instead modify the proof of the previous compact embedding and closedness results.

Cases 1–3: Theorem 3 (case 1: part 1 with the s weight equal to the constant 1 and the c weight equal to $\tilde{\mu}^2$) (case 2: part 2 with the s weight equal to 1 and the c weight equal to $\tilde{\mu}$) (case 3: part 3, with weights chosen as in case 2) implies that (cases 1 and 2: $\mathcal{W}_{m+m_0, 2, \mathbb{1}}$) (case 3: $\mathcal{C}_{m+m_0, \infty, \mathbb{1}}$) is compactly embedded in (cases 1 and 3: $\mathcal{C}_{m, \infty, \tilde{\mu}}$) (case 2: $\mathcal{W}_{m, 2, \tilde{\mu}}$). Note that both the constant weight function, $\tilde{\mu}$, and $\tilde{\mu}^2$ satisfy the local integrability assumption 6 as well as assumption 3.

By proposition 6, (cases 1 and 3: $\|\cdot\|_{m, \infty, \tilde{\mu}}$) (case 2: $\|\cdot\|_{m, 2, \tilde{\mu}}$) and (cases 1 and 3: $\|\cdot\|_{m, \infty, \tilde{\mu}, \text{ALT}}$) (case 2: $\|\cdot\|_{m, 2, \tilde{\mu}, \text{ALT}}$) are equivalent norms. Therefore (cases 1 and 2: $\mathcal{W}_{m+m_0, 2, \mathbb{1}} = \mathcal{W}_{m+m_0, 2, \mathbb{1}, \text{ALT}}$) (case 3: $\mathcal{C}_{m+m_0, \infty, \mathbb{1}, \text{ALT}}$) is compactly embedded in (cases 1 and 3: $\mathcal{C}_{m, \infty, \tilde{\mu}, \text{ALT}}$) (case 2: $\mathcal{W}_{m, 2, \tilde{\mu}, \text{ALT}}$). Lemma S1 part 1 now implies that (cases 1 and 2: $\mathcal{W}_{m+m_0, 2, \mu_s, \text{ALT}}$) (case 3: $\mathcal{C}_{m+m_0, \infty, \mu_s, \text{ALT}}$) is compactly embedded in (cases 1 and 3: $\mathcal{C}_{m, \infty, \mu_c, \text{ALT}}$) (case 2: $\mathcal{W}_{m, 2, \mu_c, \text{ALT}}$).

Case 4: The proof is similar to the proof of theorem 3. Since we have already given a detailed proof of that theorem, here we only comment on the nontrivial modifications to that proof. The numbers here refer to the steps in that proof.

1. $\Theta = \{f \in \mathcal{C}_{m+m_0, \infty, \mu_s, \nu} : \|\mu_s f\|_{m+m_0, \infty, \mathbb{1}, \nu} \leq B\}$.
2. Completeness of the function spaces under product weighting follows by completeness of the unweighted spaces.

4. This step is not necessary since, by definition of the product weighted norms, $f_n \in \Theta$ for all n implies

$\{\mu_s f_n\}$ is $\|\cdot\|_{m+m_0, \infty, \mathbb{1}, \nu}$ -bounded. In particular, this implies it is $\|\cdot\|_{m+m_0, \infty, \mathbb{1}_{\Omega_J}, \nu}$ -bounded for each J , where here

$$\|g\|_{m+m_0, \infty, \mathbb{1}_{\Omega_J}, \nu} = \|g\|_{m+m_0, \infty, \mathbb{1}_{\Omega_J}} + \max_{|\lambda|=m+m_0} \sup_{x, y \in \Omega_J, x \neq y} \frac{|\nabla^\lambda f(x) - \nabla^\lambda f(y)|}{\|x - y\|_e^\nu}.$$

Generally, in this proof indicators in the weight function placeholder denote the set over which integration or suprema are taken.

5. Apply theorem 1 part 5. Since $\{\mu_s f_n\}$ is $\|\cdot\|_{m+m_0, \infty, \mathbb{1}_{\Omega_J}, \nu}$ -bounded, it is $\|\cdot\|_{m, \infty, \mathbb{1}_{\Omega_J}}$ -relatively compact.
9. By identical calculations as before, we have

$$\|f_j - f_k\|_{m, \infty, \mu_c, \text{ALT}} \leq \|f_j - f_k\|_{m, \infty, \mu_c \mathbb{1}_{\Omega_J}, \text{ALT}} + \|f_j - f_k\|_{m, \infty, \mu_c \mathbb{1}_{\Omega_J^c}, \text{ALT}}.$$

10. For $f_j \in \Theta$ we have

$$\begin{aligned} \|f_j\|_{m, \infty, \mu_c \mathbb{1}_{\Omega_J^c}, \text{ALT}} &= \|\mu_c f_j\|_{m, \infty, \mathbb{1}_{\Omega_J^c}} \\ &= \|\mu_s \tilde{\mu} f_j\|_{m, \infty, \mathbb{1}_{\Omega_J^c}} \\ &\leq M \|\mu_s f_j\|_{m, \infty, \tilde{\mu} \mathbb{1}_{\Omega_J^c}} \\ &= M \max_{0 \leq |\lambda| \leq m} \sup_{x \in \Omega_J^c} |\nabla^\lambda (\mu_s(x) f_j(x))| \tilde{\mu}(x) \\ &\leq M \max_{0 \leq |\lambda| \leq m} \sup_{x \in \mathbb{R}^{d_x}} |\nabla^\lambda (\mu_s(x) f_j(x))| \sup_{x \in \Omega_J^c} \tilde{\mu}(x) \\ &\leq M \|\mu_s f_j\|_{m+m_0, \infty, \mathbb{1}, \nu} \sup_{x \in \Omega_J^c} \tilde{\mu}(x) \\ &\leq MB \sup_{x \in \Omega_J^c} \tilde{\mu}(x). \end{aligned}$$

The third line follows by lemma 8. The last line follows since $f_j \in \Theta$. Now since $\tilde{\mu}(x) = (1 + x'x)^{-\delta}$, $\delta > 0$, converges to zero in the tails, we can choose J large enough such that

$$\sup_{x \in \Omega_J^c} \tilde{\mu}(x) < \frac{\varepsilon}{4MB}.$$

Hence, by the triangle inequality,

$$\|f_j - f_k\|_{m, \infty, \mu_c \mathbb{1}_{\Omega_J^c}} < \frac{\varepsilon}{2}.$$

11. Since $\{\mu_s f_j^{(J)}\}$ converges in the norm $\|\cdot\|_{m,\infty,1\Omega_J}$ it is also Cauchy in that norm. Thus there is some K large enough (take $K > J$) such that

$$\|\mu_s(f_j - f_k)\|_{m,\infty,1\Omega_J} < \frac{\varepsilon}{2M}$$

for all $k, j > K$, where M is a constant given below. Hence

$$\begin{aligned} \|f_j - f_k\|_{m,\infty,\mu_c 1\Omega_J, \text{ALT}} &= \|\mu_c(f_j - f_k)\|_{m,\infty,1\Omega_J} \\ &= \|\mu_s \tilde{\mu}(f_j - f_k)\|_{m,\infty,1\Omega_J} \\ &\leq M \|\mu_s(f_j - f_k)\|_{m,\infty,\tilde{\mu} 1\Omega_J} \\ &\leq M \|\mu_s(f_j - f_k)\|_{m,\infty,1\Omega_J} \\ &< M \frac{\varepsilon}{2M} \\ &= \frac{\varepsilon}{2}. \end{aligned}$$

The third line follows by lemma 8. The fourth line follows since $\tilde{\mu}(x) = (1 + x'x)^{-\delta} \leq 1$ for all x .

□

Lemma S1. Let $(X, \|\cdot\|_X)$ and $(Y, \|\cdot\|_Y)$ be Banach spaces where $\|f\|_X < \infty$ for all $f \in X$ and $\|f\|_Y < \infty$ for all $f \in Y$. Moreover, suppose that for all $f \in X$

$$\|f\|_X = \|f\|_s$$

and for all $f \in Y$

$$\|f\|_Y = \|f\tilde{\mu}\|_c$$

where $\|\cdot\|_s$ and $\|\cdot\|_c$ are norms and $\tilde{\mu}$ is a weight function. Let $(\tilde{X}, \|\cdot\|_{\tilde{X}})$ and $(\tilde{Y}, \|\cdot\|_{\tilde{Y}})$ be Banach spaces where $\|f\|_{\tilde{X}} < \infty$ for all $f \in \tilde{X}$ and $\|f\|_{\tilde{Y}} < \infty$ for all $f \in \tilde{Y}$. Moreover, suppose that for all $f \in \tilde{X}$

$$\|f\|_{\tilde{X}} = \|f\mu_s\|_s$$

and for all $f \in \tilde{Y}$

$$\|f\|_{\tilde{Y}} = \|f\mu_s\tilde{\mu}\|_c$$

for some weight function μ_s .

1. (Compact embedding) Suppose $(X, \|\cdot\|_X)$ is compactly embedded in $(Y, \|\cdot\|_Y)$. Then $(\tilde{X}, \|\cdot\|_{\tilde{X}})$ is compactly embedded in $(\tilde{Y}, \|\cdot\|_{\tilde{Y}})$.

2. (Closedness) Suppose

$$\Omega = \{f \in X : \|f\|_X \leq B\}$$

is $\|\cdot\|_Y$ -closed. Then

$$\tilde{\Omega} = \{f \in \tilde{X} : \|f\|_{\tilde{X}} \leq B\}$$

is $\|\cdot\|_{\tilde{Y}}$ -closed.

Proof of lemma S1.

1. Let $f \in \tilde{X}$. By definition, $\|f\|_{\tilde{X}} = \|f\mu_s\|_s < \infty$. Define $h = f\mu_s$ and notice that $h \in X$. Since $(X, \|\cdot\|_X)$ is compactly embedded in $(Y, \|\cdot\|_Y)$, $X \subseteq Y$ and there exists a constant C such that $\|h\|_Y \leq C\|h\|_X$. First note that $h \in X$ implies $\|h\|_Y < \infty$ and hence $\|h\tilde{\mu}\|_c = \|f\mu_s\tilde{\mu}\|_c < \infty$. So $f \in \tilde{Y}$ and thus $\tilde{X} \subseteq \tilde{Y}$. Next, note that

$$\begin{aligned} \|h\|_Y \leq C\|h\|_X &\Leftrightarrow \|h\tilde{\mu}\|_c \leq C\|h\|_s \\ &\Leftrightarrow \|f\mu_s\tilde{\mu}\|_c \leq C\|f\mu_s\|_s \\ &\Leftrightarrow \|f\|_{\tilde{Y}} \leq C\|f\|_{\tilde{X}}. \end{aligned}$$

Next let $\{f_n\}$ be a sequence in the $\|\cdot\|_{\tilde{Y}}$ -closure of

$$\tilde{\Omega} = \{f \in \tilde{X} : \|f\|_{\tilde{X}} \leq B\} = \{f \in \tilde{X} : \|f\mu_s\|_s \leq B\}.$$

Let $h_n = f_n\mu_s$. Then by definition of the norms, h_n is a sequence in the $\|\cdot\|_Y$ -closure of

$$\Omega = \{h \in X : \|h\|_X \leq B\}.$$

Since $(X, \|\cdot\|_X)$ is compactly embedded in $(Y, \|\cdot\|_Y)$, there exists a subsequence $h_{n_j} = f_{n_j}\mu_s$, which is $\|\cdot\|_Y$ -Cauchy. That is, for any $\varepsilon > 0$, there exists an N such that $\|h_{n_j} - h_{n_k}\|_Y \leq \varepsilon$ for all $j, k > N$. But

$$\|h_{n_j} - h_{n_k}\|_Y = \|(h_{n_j} - h_{n_k})\tilde{\mu}\|_c = \|(f_{n_j} - f_{n_k})\mu_s\tilde{\mu}\|_c = \|f_{n_j} - f_{n_k}\|_{\tilde{Y}}.$$

Therefore, f_{n_j} is a subsequence of f_n which is $\|\cdot\|_{\tilde{Y}}$ -Cauchy. Since $(\tilde{Y}, \|\cdot\|_{\tilde{Y}})$ is Banach, f_j converges to a point in \tilde{Y} . Hence $(\tilde{X}, \|\cdot\|_{\tilde{X}})$ is compactly embedded in $(\tilde{Y}, \|\cdot\|_{\tilde{X}})$.

2. Let f_n be a sequence in $\tilde{\Omega}$ such that for some $f \in \tilde{X}$, $\|f_n - f\|_{\tilde{Y}} \rightarrow 0$ as $n \rightarrow \infty$. Since $f_n \in \tilde{\Omega}$ we have $\|f_n\mu_s\|_s = \|f_n\|_{\tilde{X}} \leq B$. Let $h_n = f_n\mu_s$ and $h = f\mu_s$. Since

$$\|h_n\|_X = \|h_n\|_s = \|f_n\mu_s\|_s = \|f_n\|_{\tilde{X}} \leq B$$

we have $h_n \in \Omega$. Moreover,

$$\|h_n - h\|_Y = \|(h_n - h)\tilde{\mu}\|_c = \|f_n - f\|_{\tilde{Y}} \rightarrow 0.$$

Since $\Omega = \{f \in X : \|f\|_X \leq B\}$ is $\|\cdot\|_Y$ -closed, $h \in \Omega$. That is, $f\mu_s \in \Omega$, which implies that

$$\|f\|_{\tilde{X}} = \|f\mu_s\|_X \leq B.$$

Hence $f \in \tilde{\Omega}$. So $\tilde{\Omega}$ is $\|\cdot\|_{\tilde{Y}}$ -closed. □

Proof of theorem 7 (Compact embedding for weighted norms on bounded domains). The proof is similar to the proof of theorem 3. Since we have already given a detailed proof of that theorem, here we only comment on the nontrivial modifications to that proof. The numbers here refer to the steps in that proof.

2. For case 1, $\Omega_1 = \dots = \Omega_k = \mathcal{D}$ and $\Omega^{(0)} = \mathcal{D}$ when applying Rodríguez et al. (2004).
3. We use the following more general domain truncation: Let $\{\Omega_J\}$ be a sequence of open subsets of \mathcal{D} such that
 - (a) $\Omega_J \subseteq \Omega_{J+1}$ for any J ,
 - (b) $\bigcup_{J=1}^{\infty} \Omega_J = \mathcal{D}$, and
 - (c) The closure of Ω_J does not contain the boundary of the closure of \mathcal{D} for any J . That is, $\text{Boundary}(\overline{\mathcal{D}}) \cap \overline{\Omega}_J = \emptyset$ for all J .

Roughly speaking, the sets Ω_J are converging to \mathcal{D} from the inside. They do this in such a way that for any J , the boundary points of $\overline{\mathcal{D}}$ are well separated from Ω_J .

The rest of the steps go through with very minor modifications. □

G Proofs of closedness theorems

Proof of theorem 2 (Closedness for bounded domains). For this proof we let $d_x = 1$ to simplify the notation. All arguments generalize to $d_x > 1$.

1. We want to show that the $\|\cdot\|_s = \|\cdot\|_{m+m_0,2}$ -ball Θ is $\|\cdot\|_c = \|\cdot\|_{m,\infty}$ -closed. $(\mathcal{W}_{m+m_0,2}, \|\cdot\|_{m+m_0,2})$ is compactly embedded in $(\mathcal{W}_{m,2}, \|\cdot\|_{m,2})$ by part 2 of theorem 1, which applies since we assumed \mathcal{D} satisfies the cone condition and $m_0 > d_x/2$. Lemma A.1 in Santos (2012) (reproduced in the main paper's appendix on page 2 for convenience) then implies that that the $\|\cdot\|_{m+m_0,2}$ -ball Θ is $\|\cdot\|_{m,2}$ -closed, because the Sobolev L_2 spaces are separable Hilbert spaces (theorem 3.6 of Adams and Fournier 2003). Finally, since $\|\cdot\|_{m,2} \leq \|\cdot\|_{m,\infty}$ corollary 1 implies that Θ is $\|\cdot\|_{m,\infty}$ -closed.
2. We want to show that the $\|\cdot\|_s = \|\cdot\|_{m+m_0,2}$ -ball Θ is $\|\cdot\|_c = \|\cdot\|_{m,2}$ -closed. We already showed this in the proof of part 1.

3. We want to show that the $\|\cdot\|_s = \|\cdot\|_{m+m_0, \infty}$ -ball Θ is not $\|\cdot\|_c = \|\cdot\|_{m, \infty}$ -closed. Consider the case $m = 0$ and $m_0 = 1$, so that Θ is the set of continuously differentiable functions whose levels and first derivatives are uniformly bounded by B . We will show that this set is not closed in the ordinary sup-norm $\|\cdot\|_{0, \infty}$.

Suppose $\mathcal{D} = (-1, 1)$. Define

$$g_k(x) = \sqrt{x^2 + 1/k}.$$

for integers $k \geq 1$. These are smooth approximations to the absolute value function: For each $x \in \mathcal{D}$, $g_k(x) \rightarrow \sqrt{x^2} = |x|$ as $k \rightarrow \infty$. g_k is continuous and differentiable, with first derivative

$$\begin{aligned} g'_k(x) &= \frac{1}{2}(x^2 + 1/k)^{-1/2} \cdot 2x \\ &= \frac{x}{\sqrt{x^2 + 1/k}}. \end{aligned}$$

So

$$|g'_k(x)| \leq \frac{|x|}{\sqrt{x^2 + 1/k}} \leq \frac{|x|}{\sqrt{x^2}} = 1$$

for all k . Also,

$$|g_k(x)| = \sqrt{x^2 + 1/k} \leq \sqrt{1 + 1/k} \leq \sqrt{1 + 1} = \sqrt{2}$$

for all k . Hence $g_k \in \Theta = \{f \in \mathcal{C}_1(\mathcal{D}) : \|f\|_{1, \infty} \leq B\}$ for each k , where $B = 1 + \sqrt{2}$. But, letting $f(x) = |x|$,

$$\|g_k - f\|_{0, \infty} = \sup_{x \in \mathcal{D}} |g_k(x) - f(x)| \rightarrow 0$$

as $k \rightarrow \infty$. Since f is not differentiable at 0, $f \notin \Theta$. This implies that Θ is not closed under $\|\cdot\|_{0, \infty}$.

4. We want to show that the $\|\cdot\|_s = \|\cdot\|_{m+m_0, \infty}$ -ball Θ is not $\|\cdot\|_c = \|\cdot\|_{m, 2}$ -closed. The same counterexample from part 4 applies here as well. Letting $m = 0$ and $m_0 = 1$, we will show that the $\|\cdot\|_{1, \infty}$ -ball Θ is not closed in the ordinary L_2 norm $\|\cdot\|_{0, 2}$. From part 4, we constructed a sequence g_k in Θ such that

$$\|g_k - f\|_{0, \infty} \rightarrow 0$$

as $k \rightarrow \infty$, for $f \notin \Theta$. Convergence in $\|\cdot\|_{0, \infty}$ implies convergence in $\|\cdot\|_{0, 2}$ and hence

$$\|g_k - f\|_{0, 2} \rightarrow 0$$

as $k \rightarrow \infty$. Therefore Θ is not closed under $\|f\|_{0, 2}$.

5. We want to show that $\|\cdot\|_{m+m_0, \infty, \mathbb{1}, \nu}$ -balls are $\|\cdot\|_{m, \infty}$ -closed, where $m_0 \geq 0$. Since $\|\cdot\|_{0, \infty} \leq \|\cdot\|_{m, \infty}$, corollary 1 shows that it is sufficient to prove the result for $m = 0$. That is, it is

sufficient to prove that the $\|\cdot\|_{m_0, \infty, 1, \nu}$ -ball

$$\Theta_{m_0} \equiv \{f \in \mathcal{C}_{m_0, \infty, 1, \nu} : \|f\|_{m_0, \infty, 1, \nu} \leq B\}$$

is $\|\cdot\|_{0, \infty}$ -closed, for all $m_0 \geq 0$. We proceed by induction on m_0 .

Step 1 (Base Case): Let $m_0 = 0$. We want to show that Θ_0 is $\|\cdot\|_{0, \infty}$ -closed, so we will show that its complement $\Theta_0^c = \mathcal{C}_{0, \infty} \setminus \Theta_0$ is $\|\cdot\|_{0, \infty}$ -open. That is, for any $f \in \Theta_0^c$ there exists an $\varepsilon > 0$ such that

$$\{g \in \mathcal{C}_{0, \infty} : \|f - g\|_{0, \infty} \leq \varepsilon\} \subseteq \Theta_0^c.$$

So take an arbitrary $f \in \Theta_0^c$. Since f is outside the Hölder ball Θ_0 , its Hölder norm is larger than B ,

$$\sup_{x \in \mathcal{D}} |f(x)| + \sup_{x_1, x_2 \in \mathcal{D}, x_1 \neq x_2} \frac{|f(x_1) - f(x_2)|}{|x_1 - x_2|^\nu} > B.$$

Hence there exist points $\bar{x}, \bar{x}_1, \bar{x}_2$ in the Euclidean closure of \mathcal{D} with $\bar{x}_1 \neq \bar{x}_2$ such that

$$|f(\bar{x})| + \frac{|f(\bar{x}_1) - f(\bar{x}_2)|}{|\bar{x}_1 - \bar{x}_2|^\nu} > B.$$

Define

$$\delta = |f(\bar{x})| + \frac{|f(\bar{x}_1) - f(\bar{x}_2)|}{|\bar{x}_1 - \bar{x}_2|^\nu} - B > 0.$$

Our goal is find a $\|\cdot\|_{0, \infty}$ -ball around f with some positive radius ε such that all functions g in that ball are also not in the Hölder ball Θ_0 . So we need these functions g to have a large Hölder norm (larger than B). Let's examine that. For all $g \in \mathcal{C}_{0, \infty}$,

$$\begin{aligned} \|g\|_{0, \infty, 1, \nu} &= \sup_{x \in \mathcal{D}} |g(x)| + \sup_{x_1, x_2 \in \mathcal{D}, x_1 \neq x_2} \frac{|g(x_1) - g(x_2)|}{|x_1 - x_2|^\nu} \\ &\geq |g(\bar{x})| + \frac{|g(\bar{x}_1) - g(\bar{x}_2)|}{|\bar{x}_1 - \bar{x}_2|^\nu} \\ &\geq |f(\bar{x})| - |f(\bar{x}) - g(\bar{x})| + \frac{|g(\bar{x}_1) - g(\bar{x}_2)|}{|\bar{x}_1 - \bar{x}_2|^\nu} \\ &= |f(\bar{x})| - |f(\bar{x}) - g(\bar{x})| \\ &\quad + \frac{|f(\bar{x}_1) - f(\bar{x}_2)|}{|\bar{x}_1 - \bar{x}_2|^\nu} - \frac{|f(\bar{x}_1) - f(\bar{x}_2)|}{|\bar{x}_1 - \bar{x}_2|^\nu} + \frac{|g(\bar{x}_1) - g(\bar{x}_2)|}{|\bar{x}_1 - \bar{x}_2|^\nu} \\ &\geq |f(\bar{x})| - |f(\bar{x}) - g(\bar{x})| \\ &\quad + \frac{|f(\bar{x}_1) - f(\bar{x}_2)|}{|\bar{x}_1 - \bar{x}_2|^\nu} - \frac{|(f(\bar{x}_1) - g(\bar{x}_1)) - (f(\bar{x}_2) - g(\bar{x}_2))|}{|\bar{x}_1 - \bar{x}_2|^\nu} \\ &= B + \delta - \left(|f(\bar{x}) - g(\bar{x})| + \frac{|(f(\bar{x}_1) - g(\bar{x}_1)) - (f(\bar{x}_2) - g(\bar{x}_2))|}{|\bar{x}_1 - \bar{x}_2|^\nu} \right). \end{aligned}$$

The third and fifth lines follow by the reverse triangle inequality. The last line follows by the

definition of δ . If we can make this last piece in parentheses small enough, we'll be done. For any $\varepsilon > 0$,

$$g \in \{g \in \mathcal{C}_{0,\infty} : \|f - g\|_{0,\infty} \leq \varepsilon\}$$

implies

$$|f(\bar{x}) - g(\bar{x})| + \frac{|(f(\bar{x}_1) - g(\bar{x}_1)) - (f(\bar{x}_2) - g(\bar{x}_2))|}{|\bar{x}_1 - \bar{x}_2|^\nu} \leq \varepsilon + \frac{2\varepsilon}{|\bar{x}_1 - \bar{x}_2|^\nu}$$

by the triangle inequality. So suppose we choose ε so that

$$\varepsilon + \frac{2\varepsilon}{|\bar{x}_1 - \bar{x}_2|^\nu} \leq \frac{\delta}{2}.$$

Note that this choice of ε depends on the particular $f \in \Theta_0^c$ chosen at the beginning, via δ and \bar{x}_1 and \bar{x}_2 . Then for all $g \in \mathcal{C}_{0,\infty}$ with $\|f - g\|_{0,\infty} \leq \varepsilon$ we have

$$\begin{aligned} \|g\|_{0,\infty,1,\nu} &\geq B + \delta - \frac{\delta}{2} \\ &= B + \frac{\delta}{2} \\ &> B. \end{aligned}$$

Hence $g \in \Theta_0^c$ for all such g . Thus Θ_0^c is $\|\cdot\|_{0,\infty}$ -open and hence Θ_0 is $\|\cdot\|_{0,\infty}$ -closed.

Step 2 (Induction Step): Next we suppose that Θ_{m_0} is $\|\cdot\|_{0,\infty}$ -closed for some integer $m_0 \geq 0$. We will show that this implies Θ_{m_0+1} is $\|\cdot\|_{0,\infty}$ -closed.

Since Θ_{m_0} is $\|\cdot\|_{0,\infty}$ -closed, we have that for all f in $\Theta_{m_0}^c = \mathcal{C}_{0,\infty} \setminus \Theta_{m_0}$ there exists an $\varepsilon > 0$ such that for all $g \in \mathcal{C}_{0,\infty}$ with

$$\|f - g\|_{0,\infty} \leq \varepsilon,$$

it holds that $g \in \Theta_{m_0}^c$. As in the base case, we will show that $\Theta_{m_0+1}^c$ is $\|\cdot\|_{0,\infty}$ -open. So take an arbitrary $f \in \Theta_{m_0+1}^c$. We will show that there exists an $\varepsilon > 0$ such that for all $g \in \mathcal{C}_{0,\infty}$ with $\|f - g\|_{0,\infty} \leq \varepsilon$ we have $g \in \Theta_{m_0+1}^c$. We have to consider several cases, depending on the properties of the f we're given. First, $\Theta_{m_0+1} \subsetneq \Theta_{m_0}$ implies

$$\Theta_{m_0}^c \subsetneq \Theta_{m_0+1}^c.$$

So it might be the case that $f \in \Theta_{m_0}^c$. This is case (a) below. Moreover, it is possible that $f \in \Theta_{m_0+1}^c$ but $f \notin \Theta_{m_0}^c$. This case could occur for several reasons. It might be that $f \in \mathcal{C}_{m_0+1,\infty,1,\nu}$, so $\|f\|_{m_0+1,\infty,1,\nu} \leq D$ for some constant $D < \infty$, but that this norm, while finite, is still too big:

$$\|f\|_{m_0+1,\infty,1,\nu} > B.$$

This is case (b) below. Another possibility is that $f \notin \mathcal{C}_{m_0+1,\infty,1,\nu}$. But $f \notin \Theta_{m_0}^c$, $f \in \Theta_{m_0}$ and hence its m_0 'th derivative exists and is Hölder continuous. So there are three reasons

why $f \notin \mathcal{C}_{m_0+1,\infty,1,\nu}$ could occur: Either the $(m_0 + 1)$ 'th derivative does not exist (case (c) below), the $(m_0 + 1)$ 'th derivative exists but is not $\|\cdot\|_{0,\infty}$ -bounded (i.e., the first piece of the Hölder norm $\|f\|_{m_0+1,\infty,1,\nu}$ is infinite) (case (d) below), or the $(m_0 + 1)$ 'th derivative exists and is $\|\cdot\|_{0,\infty}$ -bounded, but is not Hölder continuous (i.e., the first piece of the Hölder norm $\|f\|_{m_0+1,\infty,1,\nu}$ is finite, but the second piece is infinite) (case (e) below).

- (a) Suppose $f \in \Theta_{m_0}^c$. But we already know from the induction assumption that $\Theta_{m_0}^c$ is open. Hence there exists an $\varepsilon > 0$ such that for all $g \in \mathcal{C}_{0,\infty}$ with $\|f - g\|_{0,\infty} \leq \varepsilon$ it holds that $g \in \Theta_{m_0}^c \subsetneq \Theta_{m_0+1}^c$.
- (b) Suppose $f \notin \Theta_{m_0}^c$ and $f \in \mathcal{C}_{m_0+1,\infty,1,\nu}$ with

$$B < \|f\|_{m_0+1,\infty,1,\nu} \leq D$$

for some constant $D < \infty$. Since $f \notin \Theta_{m_0}^c$, $f \in \Theta_{m_0}$ and hence

$$\|f\|_{m_0,\infty,1,\nu} \leq B.$$

Let $g \in \mathcal{C}_{0,\infty}$ be such that $\|f - g\|_{0,\infty} \leq \varepsilon$. Remember that our goal is to find an $\varepsilon > 0$ such that all of these g are in $\Theta_{m_0+1}^c$. Regardless of the value of ε , if $g \notin \mathcal{C}_{m_0+1,\infty,1,\nu}$ (in which case $g \notin \Theta_{m_0+1}$ and so $g \in \Theta_{m_0+1}^c$) or if $\|g\|_{m_0+1,\infty,1,\nu} \geq C$ for some finite constant $C > B$, then $g \in \Theta_{m_0+1}^c$. So suppose that $g \in \mathcal{C}_{m_0+1,\infty,1,\nu}$ and

$$\|g\|_{m_0+1,\infty,1,\nu} \leq C.$$

We will show that although this norm is smaller than C , it is still larger than B . For each $x \in \mathcal{D}$ and $\delta > 0$ with $x + \delta \in \mathcal{D}$,² the mean value theorem implies that there exists an $x_g \in [x, x + \delta]$ such that

$$g'(x_g) = \frac{g(x + \delta) - g(x)}{\delta}$$

and hence

$$\begin{aligned} g'(x) &= g'(x_g) + (g'(x) - g'(x_g)) \\ &= \frac{g(x + \delta) - g(x)}{\delta} + (g'(x) - g'(x_g)). \end{aligned}$$

Note that g is differentiable because $g \in \mathcal{C}_{m_0+1,\infty,1,\nu}$. Likewise, there exists an $x_f \in [x, x + \delta]$ such that

$$f'(x) = \frac{f(x + \delta) - f(x)}{\delta} + (f'(x) - f'(x_f)).$$

²The cone condition implies that there exists a single $\delta > 0$ such that, for all $x \in \mathcal{D}$, at least one of $x + \delta \in \mathcal{D}$ or $x - \delta \in \mathcal{D}$ holds.

It follows that

$$\begin{aligned}
& \|f' - g'\|_{0,\infty} \\
&= \sup_{x \in \mathcal{D}} |f'(x) - g'(x)| \\
&= \sup_{x \in \mathcal{D}} \left| \left(\frac{f(x+\delta) - f(x)}{\delta} + (f'(x) - f'(x_f)) \right) - \left(\frac{g(x+\delta) - g(x)}{\delta} + (g'(x) - g'(x_g)) \right) \right| \\
&= \sup_{x \in \mathcal{D}} \left| \frac{f(x+\delta) - g(x+\delta)}{\delta} - \frac{f(x) - g(x)}{\delta} + (f'(x) - f'(x_f)) + (g'(x) - g'(x_g)) \right| \\
&\leq \sup_{x \in \mathcal{D}} \left(\frac{|f(x+\delta) - g(x+\delta)|}{\delta} + \frac{|f(x) - g(x)|}{\delta} + |f'(x) - f'(x_f)| + |g'(x) - g'(x_g)| \right) \\
&\leq \frac{2\varepsilon}{\delta} + D\delta^\nu + C\delta^\nu
\end{aligned}$$

The fourth line follows by the triangle inequality. The last line by $\|f - g\|_{0,\infty} \leq \varepsilon$, $x_f \in [x, x + \delta]$, $x_g \in [x, x + \delta]$, and since f' and g' are both Hölder continuous with Hölder constants D and C , respectively (which follows because $\|f\|_{m_0+1,\infty,1,\nu} \leq D$ and $\|g\|_{m_0+1,\infty,1,\nu} \leq C$).

Let $\varepsilon_1 > 0$ be arbitrary. Choose $\delta > 0$ such that $D\delta^\nu \leq \varepsilon_1/3$ and $C\delta^\nu \leq \varepsilon_1/3$. After choosing δ , choose ε such that $2\varepsilon/\delta \leq \varepsilon_1/3$. Thus

$$\|f' - g'\|_{0,\infty} \leq \varepsilon_1.$$

We have shown that if the first derivatives of f and g are Hölder continuous, we can make the derivatives for all g with $\|f - g\|_{0,\infty} \leq \varepsilon$ arbitrarily close to the derivative of f by choosing ε small enough. An analogous argument shows that if $\|f' - g'\|_{0,\infty} \leq \varepsilon_1$ and if the second derivatives are Hölder continuous, then we can make the second derivatives arbitrarily close. Applying this argument recursively to higher order derivative shows that for any $\varepsilon_{m_0+1} > 0$, we can pick an $\varepsilon > 0$ such that for all g with $\|g\|_{m_0+1,\infty,1,\nu} \leq C$ and $\|f - g\|_{0,\infty} \leq \varepsilon$,

$$\|\nabla^{m_0+1} f - \nabla^{m_0+1} g\|_{0,\infty} \leq \varepsilon_{m_0+1}.$$

Our argument from the base case (step 1) now implies that if ε_{m_0+1} is small enough, then $\|g\|_{m_0+1,\infty,1,\nu} > B$ for all $g \in \mathcal{C}_{0,\infty}$ with $\|f - g\|_{0,\infty} \leq \varepsilon$. Hence $g \in \Theta_{m_0+1}^c$. Note that we use $\|f\|_{m_0+1,\infty,1,\nu} > B$ when applying the base case argument.

- (c) Suppose that for some $\bar{x} \in \mathcal{D}$, $\nabla^{m_0+1} f(\bar{x})$ does not exist. Then $f \notin \mathcal{C}_{m_0+1,\infty,1,\nu}^c$. But since $f \notin \Theta_{m_0}^c$, we know that the m_0 'th derivative of f exists and is Hölder continuous. As in case (b), take $g \in \mathcal{C}_{0,\infty}$ such that $\|f - g\|_{0,\infty} \leq \varepsilon$ and suppose that $g \in \mathcal{C}_{m_0+1,\infty,1,\nu}$ with $\|g\|_{m_0+1,\infty,1,\nu} \leq C$ for $C > B$ (remember from part (b) that otherwise we know $g \in \Theta_{m_0+1}^c$ already). Since the m_0 'th derivative of f exists and is Hölder continuous, we know that the only way for the derivative $\nabla^{m_0+1} f(\bar{x})$ to not exist is if it has a kink—its right hand side derivative does not exist, its left hand side derivative does not exist, or

both exist but are not equal. So we consider each of these three cases separately.

i. Suppose the right hand side derivative of $\nabla^{m_0} f$ at \bar{x} does not exist. That is,

$$\lim_{h \searrow 0} \frac{\nabla^{m_0} f(\bar{x} + h) - \nabla^{m_0} f(\bar{x})}{h}$$

does not exist. Then there exists a $\delta > 0$ such that for any $\eta > 0$ we can find an h with $0 < h < \eta$ and

$$\left| \frac{\nabla^{m_0} f(\bar{x} + h) - \nabla^{m_0} f(\bar{x})}{h} - \nabla^{m_0+1} g(\bar{x}) \right| > \delta.$$

If such a δ did not exist, then

$$\lim_{h \searrow 0} \frac{\nabla^{m_0} f(\bar{x} + h) - \nabla^{m_0} f(\bar{x})}{h} = \nabla^{m_0+1} g(\bar{x})$$

by definition of the limit. For such a fixed h , we have

$$\begin{aligned} \delta &< \left| \frac{\nabla^{m_0} f(\bar{x} + h) - \nabla^{m_0} f(\bar{x})}{h} - \nabla^{m_0+1} g(\bar{x}) \right| \\ &\leq \left| \frac{\nabla^{m_0} f(\bar{x} + h) - \nabla^{m_0} g(\bar{x} + h) + \nabla^{m_0} g(\bar{x}) - \nabla^{m_0} f(\bar{x})}{h} \right| \\ &\quad + \left| \frac{\nabla^{m_0} g(\bar{x} + h) - \nabla^{m_0} g(\bar{x})}{h} - \nabla^{m_0+1} g(\bar{x}) \right| \\ &\leq \left| \frac{\nabla^{m_0} f(\bar{x} + h) - \nabla^{m_0} g(\bar{x} + h) + \nabla^{m_0} g(\bar{x}) - \nabla^{m_0} f(\bar{x})}{h} \right| \\ &\quad + |\nabla^{m_0+1} g(\tilde{x}) - \nabla^{m_0+1} g(\bar{x})| \\ &\leq \left| \frac{\nabla^{m_0} f(\bar{x} + h) - \nabla^{m_0} g(\bar{x} + h) + \nabla^{m_0} g(\bar{x}) - \nabla^{m_0} f(\bar{x})}{h} \right| + Ch^\nu. \end{aligned}$$

The second line follows by the triangle inequality. The third line by the mean value theorem, since $\nabla^{m_0} g$ is differentiable, and here $\tilde{x} \in [\bar{x}, \bar{x} + h]$. The fourth line follows since $\nabla^{m_0+1} g$ is Hölder continuous with constant C , and since $\tilde{x} \in [\bar{x}, \bar{x} + h]$ so that $\|\tilde{x} - \bar{x}\| \leq h$. Now choose h small enough such that $Ch^\nu \leq \delta/2$. For this fixed h , pick ε small enough such that

$$\|\nabla^{m_0} f - \nabla^{m_0} g\|_{0,\infty} \leq \frac{\delta h}{4}.$$

Then

$$\delta < \left| \frac{\nabla^{m_0} f(\bar{x} + h) - \nabla^{m_0} f(\bar{x})}{h} - \nabla^{m_0+1} g(\bar{x}) \right| \leq \delta,$$

a contraction.

ii. Suppose the left hand side derivative of $\nabla^{m_0} f$ at \bar{x} does not exist. That is,

$$\lim_{h \searrow 0} \frac{\nabla^{m_0} f(\bar{x}) - \nabla^{m_0} f(\bar{x} - h)}{h}$$

does not exist. This case proceeds analogously to the previous case.

iii. Both the left hand and right hand side derivatives of $\nabla^{m_0} f$ at \bar{x} exist, but they are not equal:

$$\lim_{h \searrow 0} \frac{\nabla^{m_0} f(\bar{x} + h) - \nabla^{m_0} f(\bar{x})}{h} \neq \lim_{h \searrow 0} \frac{\nabla^{m_0} f(\bar{x}) - \nabla^{m_0} f(\bar{x} - h)}{h}.$$

Considering the distance between the right hand side and left hand side secant lines, for any $h > 0$ such that $[\bar{x} - h, \bar{x} + h] \subseteq \mathcal{D}$, we obtain

$$\begin{aligned} & \left| \left(\frac{\nabla^{m_0} f(\bar{x} + h) - \nabla^{m_0} f(\bar{x})}{h} \right) - \left(\frac{\nabla^{m_0} f(\bar{x}) - \nabla^{m_0} f(\bar{x} - h)}{h} \right) \right| \\ & \leq 4 \frac{\varepsilon_{m_0}}{h} + \left| \left(\frac{\nabla^{m_0} g(\bar{x} + h) - \nabla^{m_0} g(\bar{x})}{h} \right) - \left(\frac{\nabla^{m_0} g(\bar{x}) - \nabla^{m_0} g(\bar{x} - h)}{h} \right) \right| \\ & = 4 \frac{\varepsilon_{m_0}}{h} + |(\nabla^{m_0+1} g(\tilde{x}_1) - \nabla^{m_0+1} g(\tilde{x}_2))| \\ & \leq 4 \frac{\varepsilon_{m_0}}{h} + C(2h)^\nu. \end{aligned}$$

For the first line, we used the triangle inequality plus the fact that for any $\varepsilon_{m_0} > 0$, there exists an $\varepsilon > 0$ not depending on g such that $\|f - g\|_{0,\infty} \leq \varepsilon$ implies

$$\|\nabla^{m_0} f - \nabla^{m_0} g\|_{0,\infty} \leq \varepsilon_{m_0}.$$

This follows from our argument in part (b), since $\nabla^{m_0} f$ and $\nabla^{m_0} g$ are Hölder continuous.

In the second line, we used the mean value theorem, since $g \in \mathcal{C}_{m_0+1,\infty,1,\nu}$, where $\tilde{x}_1 \in [\bar{x}, \bar{x} + h]$ and $\tilde{x}_2 \in [\bar{x} - h, \bar{x}]$. In the third line we used Hölder continuity of $\nabla^{m_0+1} g$ since $\|g\|_{m_0+1,\infty,1,\nu} \leq C$, plus the fact that $|\tilde{x}_1 - \tilde{x}_2| \leq 2h$.

Since

$$\lim_{h \searrow 0} \frac{\nabla^{m_0} f(\bar{x} + h) - \nabla^{m_0} f(\bar{x})}{h} \neq \lim_{h \searrow 0} \frac{\nabla^{m_0} f(\bar{x}) - \nabla^{m_0} f(\bar{x} - h)}{h}$$

there exists a $\delta > 0$ such that for an arbitrarily small h

$$\left| \left(\frac{\nabla^{m_0} f(\bar{x} + h) - \nabla^{m_0} f(\bar{x})}{h} \right) - \left(\frac{\nabla^{m_0} f(\bar{x}) - \nabla^{m_0} f(\bar{x} - h)}{h} \right) \right| > \delta.$$

Choose h such that $C(2h)^\nu \leq \delta/2$. Then for this fixed h , pick ε small enough such

that $4\varepsilon_{m_0}/h \leq \delta/2$. Then

$$\delta < \left| \left(\frac{\nabla^{m_0} f(\bar{x} + h) - \nabla^{m_0} f(\bar{x})}{h} \right) - \left(\frac{\nabla^{m_0} f(\bar{x}) - \nabla^{m_0} f(\bar{x} - h)}{h} \right) \right| \leq \delta,$$

a contraction.

In all three cases where $\nabla^{m_0+1} f(\bar{x})$ does not exist, we have derived a contradiction. Hence there does not exist a $g \in \mathcal{C}_{0,\infty}$ with $\|g\|_{m_0+1,\infty,1,\nu} \leq C$ and $\|f - g\|_{0,\infty} \leq \varepsilon$. This implies that for all $g \in \mathcal{C}_{0,\infty}$ with $\|f - g\|_{0,\infty} \leq \varepsilon$ it holds that $g \in \Theta_{m_0+1}^c$.

(d) Suppose $\nabla^{m_0+1} f(x)$ exists for all $x \in \mathcal{D}$ but

$$\sup_{x \in \mathcal{D}} |\nabla^{m_0+1} f(x)| = \infty.$$

For example, this happens with $f(x) = \sqrt{x}$ when $\mathcal{D} = (0, 1)$ and $m_0 = 0$. Then there exists a $\bar{x} \in \mathcal{D}$ such that

$$C < |\nabla^{m_0+1} f(\bar{x})| < \infty$$

for some constant $C > B$. Thus, for all $\|g\|_{m_0+1,\infty,1,\nu} \leq C$,

$$\begin{aligned} |\nabla^{m_0+1} g(\bar{x})| &\geq |\nabla^{m_0+1} f(\bar{x})| - |\nabla^{m_0+1} g(\bar{x}) - \nabla^{m_0+1} f(\bar{x})| \\ &= |\nabla^{m_0+1} f(\bar{x})| - \left| \lim_{h \rightarrow 0} \frac{\nabla^{m_0} g(\bar{x} + h) - \nabla^{m_0} g(\bar{x})}{h} - \lim_{h \rightarrow 0} \frac{\nabla^{m_0} f(\bar{x} + h) - \nabla^{m_0} f(\bar{x})}{h} \right| \\ &= |\nabla^{m_0+1} f(\bar{x})| - \lim_{h \rightarrow 0} \left| \frac{\nabla^{m_0} g(\bar{x} + h) - \nabla^{m_0} f(\bar{x} + h)}{h} - \frac{\nabla^{m_0} g(\bar{x}) - \nabla^{m_0} f(\bar{x})}{h} \right|. \end{aligned}$$

The first line follows by the reverse triangle inequality. Since the limit in the last line exists and is finite, for any $\delta > 0$, we can find an $\bar{h} > 0$ with $[\bar{x}, \bar{x} + \bar{h}] \subseteq \mathcal{D}$ such that the difference between the limit and the term we're taking the limit of evaluated at \bar{h} is smaller than δ . Hence

$$\begin{aligned} |\nabla^{m_0+1} g(\bar{x})| &\geq |\nabla^{m_0+1} f(\bar{x})| - \left| \frac{\nabla^{m_0} g(\bar{x} + \bar{h}) - \nabla^{m_0} f(\bar{x} + \bar{h})}{\bar{h}} - \frac{\nabla^{m_0} g(\bar{x}) - \nabla^{m_0} f(\bar{x})}{\bar{h}} \right| - \delta \\ &\geq C - \delta - \left| \frac{\nabla^{m_0} g(\bar{x} + \bar{h}) - \nabla^{m_0} f(\bar{x} + \bar{h})}{\bar{h}} - \frac{\nabla^{m_0} g(\bar{x}) - \nabla^{m_0} f(\bar{x})}{\bar{h}} \right|. \end{aligned}$$

As in part (b), for any $\varepsilon_{m_0} > 0$, there is an $\varepsilon > 0$ such that $\|f - g\|_{0,\infty} \leq \varepsilon$ implies

$$\|\nabla^{m_0} f - \nabla^{m_0} g\|_{0,\infty} \leq \varepsilon_{m_0}.$$

Let ε_{m_0} such that

$$\left| \frac{\nabla^{m_0} g(\bar{x} + \bar{h}) - \nabla^{m_0} f(\bar{x} + \bar{h})}{\bar{h}} - \frac{\nabla^{m_0} g(\bar{x}) - \nabla^{m_0} f(\bar{x})}{\bar{h}} \right| \leq \delta.$$

Then

$$\begin{aligned} |\nabla^{m_0+1}g(\bar{x})| &\geq C - 2\delta \\ &> B \end{aligned}$$

where the last line follows if we choose $\delta > 0$ such that $C - 2\delta > B$, that is, $\delta < (C - B)/2$, which is possible since $C > B$. We have shown that the first piece of the Hölder norm $\|g\|_{m_0+1, \infty, 1, \nu}$ is larger than B , and so the entire norm is larger than B and hence $g \in \Theta_{m_0+1}^c$.

(e) Finally, suppose

$$\sup_{x \in \mathcal{D}} |\nabla^{m_0+1}f(x)| \leq D < \infty$$

but $\nabla^{m_0+1}f$ is not Hölder continuous:

$$\sup_{x_1, x_2 \in \mathcal{D}, x_1 \neq x_2} \frac{|\nabla^{m_0+1}f(x_1) - \nabla^{m_0+1}f(x_2)|}{|x_1 - x_2|^\nu} = \infty.$$

Again take $g \in \mathcal{C}_{0, \infty}$ such that $\|f - g\|_{0, \infty} \leq \varepsilon$ and suppose that $\|g\|_{m_0+1, \infty, 1, \nu} \leq C$ for $C > B$. Since $\nabla^{m_0+1}f$ is not Hölder continuous, there exist x_1 and x_2 in \mathcal{D} , $x_1 \neq x_2$, such that

$$\left| \frac{\nabla^{m_0+1}f(x_1) - \nabla^{m_0+1}f(x_2)}{|x_1 - x_2|^\nu} \right| > B + C.$$

Moreover, by the triangle inequality,

$$\begin{aligned} &\left| \frac{\nabla^{m_0+1}f(x_1) - \nabla^{m_0+1}f(x_2)}{|x_1 - x_2|^\nu} \right| \\ &\leq \left| \frac{\nabla^{m_0+1}g(x_1) - \nabla^{m_0+1}g(x_2)}{|x_1 - x_2|^\nu} \right| + \\ &+ \lim_{h \rightarrow 0} \left| \frac{(\nabla^{m_0}g(x_1 + h) - \nabla^{m_0}g(x_1)) - (\nabla^{m_0}f(x_1 + h) - \nabla^{m_0}f(x_1))}{h} \right| / |x_1 - x_2|^\nu \\ &+ \lim_{h \rightarrow 0} \left| \frac{(\nabla^{m_0}g(x_2 + h) - \nabla^{m_0}g(x_2)) - (\nabla^{m_0}f(x_2 + h) - \nabla^{m_0}f(x_2))}{h} \right| / |x_1 - x_2|^\nu. \end{aligned}$$

As in part (b), for any $\varepsilon_{m_0} > 0$, there is an $\varepsilon > 0$ such that $\|f - g\|_{0, \infty} \leq \varepsilon$ implies

$$\|\nabla^{m_0}f - \nabla^{m_0}g\|_{0, \infty} \leq \varepsilon_{m_0}.$$

Returning to our previous inequality, we see that since the limits on the right hand side are finite and since $\nabla^{m_0+1}g$ is Hölder continuous, for any $\delta > 0$ there is an $\bar{h} > 0$ which

does not depend on g such that

$$\begin{aligned}
& \left| \frac{\nabla^{m_0+1} f(x_1) - \nabla^{m_0+1} f(x_2)}{|x_1 - x_2|^\nu} \right| \\
& \leq \left| \frac{\nabla^{m_0+1} g(x_1) - \nabla^{m_0+1} g(x_2)}{|x_1 - x_2|^\nu} \right| \\
& + \left| \frac{(\nabla^{m_0} g(x_1 + \bar{h}) - \nabla^{m_0} g(x_1)) - (\nabla^{m_0} f(x_1 + \bar{h}) - \nabla^{m_0} f(x_1))}{\bar{h}} \right| / |x_1 - x_2|^\nu \\
& + \left| \frac{(\nabla^{m_0} g(x_2 + \bar{h}) - \nabla^{m_0} g(x_2)) - (\nabla^{m_0} f(x_2 + \bar{h}) - \nabla^{m_0} f(x_2))}{\bar{h}} \right| / |x_1 - x_2|^\nu + \delta \\
& \leq C + \frac{4\varepsilon_{m_0}}{\bar{h}|x_1 - x_2|^\nu} + \delta.
\end{aligned}$$

This is the same argument we used in part (d). In the last line we used $\|g\|_{m_0+1, \infty, 1, \nu} \leq C$, the triangle inequality, and $\|\nabla^{m_0} f - \nabla^{m_0} g\|_{0, \infty} \leq \varepsilon_{m_0}$. Choose $\delta = B/2$. Then choose ε_{m_0} small enough so that

$$\frac{4\varepsilon_0}{\bar{h}|x_1 - x_2|^\nu} < \frac{B}{2}.$$

Combining our results, we have shown

$$C + B < \left| \frac{\nabla^{m_0+1} f(x_1) - \nabla^{m_0+1} f(x_2)}{|x_1 - x_2|^\nu} \right| \leq C + B,$$

a contradiction. □

Proof of theorem 4 (Closedness under equal weightings).

1. We want to show that the $\|\cdot\|_s = \|\cdot\|_{m+m_0, 2, \mu_s}$ -ball Θ is $\|\cdot\|_c = \|\cdot\|_{m, \infty, \mu_c^{1/2}}$ -closed. Part 1 of our compact embedding result theorem 3 says that $\mathscr{W}_{m+m_0, 2, \mu_s}$ is compactly embedded in $\mathscr{C}_{m, \infty, \mu_c^{1/2}}$. Now consider the space $(\mathscr{W}_{m, 2, \mu_a}, \|\cdot\|_{m, 2, \mu_a})$ where μ_a is such that

$$\int_{\mathbb{R}^{d_x}} \frac{\mu_a(x)}{\mu_c(x)} dx \leq C_1.$$

Then for any $f \in \mathcal{C}_{m,\infty,\mu_c}^{1/2}$,

$$\begin{aligned}
\|f\|_{m,2,\mu_a}^2 &= \sum_{0 \leq |\lambda| \leq m} \int_{\mathbb{R}^{d_x}} |\nabla^\lambda f(x)|^2 \mu_a(x) dx \\
&= \sum_{0 \leq |\lambda| \leq m} \int_{\mathbb{R}^{d_x}} |\nabla^\lambda f(x)|^2 \mu_c(x) \frac{\mu_a(x)}{\mu_c(x)} dx \\
&\leq C \|f\|_{m,\infty,\mu_c}^2 \int_{\mathbb{R}^{d_x}} \frac{\mu_a(x)}{\mu_c(x)} dx \\
&\leq CC_1 \|f\|_{m,\infty,\mu_c}^{1/2}.
\end{aligned}$$

Hence

$$\mathcal{C}_{m,\infty,\mu_c}^{1/2} \subseteq \mathcal{W}_{m,2,\mu_a}.$$

But we also know that $\mathcal{W}_{m+m_0,2,\mu_s}$ is compactly embedding in $\mathcal{C}_{m,\infty,\mu_c}^{1/2}$. Therefore, by lemma 4, $\mathcal{W}_{m+m_0,2,\mu_s}$ is compactly embedded in $\mathcal{W}_{m,2,\mu_a}$. Both of these are separable Hilbert spaces by arguments as in the proof of theorem 3.6 in Kufner (1980), which is analogous to Adams and Fournier (2003) theorem 3.6. Hence lemma A.1 of Santos (2012) implies that Θ is $\|\cdot\|_{m,2,\mu_a}$ -closed. But now lemma 2 and the inequality $\|\cdot\|_{m,2,\mu_a} \leq (CC_1)^{1/2} \|\cdot\|_{m,\infty,\mu_c}^{1/2}$ imply that Θ is $\|\cdot\|_{m,\infty,\mu_c}^{1/2}$ -closed.

2. We want to show that the $\|\cdot\|_s = \|\cdot\|_{m+m_0,2,\mu_s}$ -ball Θ is $\|\cdot\|_c = \|\cdot\|_{m,2,\mu_c}$ -closed. Part 2 of our compact embedding result theorem 3 says that $\mathcal{W}_{m+m_0,2,\mu_s}$ is compactly embedded in $\mathcal{W}_{m,2,\mu_c}$. Both of these are separable Hilbert spaces, as discussed in the previous part. Hence lemma A.1. of Santos (2012) implies that Θ is $\|\cdot\|_{m,2,\mu_c}$ -closed.
3. We want to show that the $\|\cdot\|_s = \|\cdot\|_{m+m_0,\infty,\mu_s}$ -ball Θ is not $\|\cdot\|_c = \|\cdot\|_{m,\infty,\mu_c}$ -closed. The same counterexample from the proof of part 3 of theorem 2 can be adapted here as well, by smoothly extending its domain definition to $\mathcal{D} = \mathbb{R}$.
4. We want to show that the $\|\cdot\|_s = \|\cdot\|_{m+m_0,\infty,\mu_s}$ -ball Θ is not $\|\cdot\|_c = \|\cdot\|_{m,2,\mu_c}$ -closed. As in the previous part, this can be shown by extending the same counterexample from theorem 2.

□

Proof of theorem 6 (Closedness under product weightings). Cases 1 and 2. This follows exactly as in the proof of theorem 5, except we apply theorem 4 and then lemma S1 part 2

Case 3. As in theorem 4, we can adapt the counterexample from theorem 2 by smoothly extending its domain to $\mathcal{D} = \mathbb{R}$.

Case 4. Assume $d_x = 1$ for simplicity. This proof is a close modification to the corresponding proof of theorem 2 for bounded domains. As in that proof, it suffices to prove the result for $m = 0$. For any $g \in \mathcal{C}_{m_0,\infty,\mu_s,\nu}$ define $g_s(x) = \mu_s(x)g(x)$ and $g_c(x) = \mu_c(x)g(x)$. We want to prove that

$$\Theta_{m_0} \equiv \{g \in \mathcal{C}_{m_0,\infty,\mu_s,\nu} : \|g\|_{m_0,\infty,\mu_s,\nu} \leq B\}$$

is $\|\cdot\|_{m_0, \infty, \mu_c}$ -closed, for all $m_0 \geq 0$. We proceed by induction on m_0 .

Step 1 (Base Case): Let $m_0 = 0$. We want to show that Θ_0 is $\|\cdot\|_{0, \infty, \mu_c}$ -closed, so we will show that its complement $\Theta_0^c = \mathcal{C}_{0, \infty, \mu_c} \setminus \Theta_0$ is $\|\cdot\|_{0, \infty, \mu_c}$ -open. So take an arbitrary $f \in \Theta_0^c$. We will show that there exists an $\varepsilon > 0$ such that

$$\{g \in \mathcal{C}_{0, \infty, \mu_c} : \|f - g\|_{0, \infty, \mu_c} \leq \varepsilon\} \subseteq \Theta_0^c.$$

Since f is outside the weighted Hölder ball Θ_0 , its weighted Hölder norm is larger than B ,

$$\sup_{x \in \mathbb{R}} |f_s(x)| + \sup_{x_1, x_2 \in \mathbb{R}} \frac{|f_s(x_1) - f_s(x_2)|}{|x_1 - x_2|^\nu} > B.$$

Hence there exist points $\bar{x}, \bar{x}_1, \bar{x}_2 \in \mathbb{R}$ with $\bar{x}_1 \neq \bar{x}_2$ such that

$$|f_s(\bar{x})| + \frac{|f_s(\bar{x}_1) - f_s(\bar{x}_2)|}{|\bar{x}_1 - \bar{x}_2|^\nu} > B.$$

Define

$$\delta = |f_s(\bar{x})| + \frac{|f_s(\bar{x}_1) - f_s(\bar{x}_2)|}{|\bar{x}_1 - \bar{x}_2|^\nu} - B > 0.$$

Next, for all $g \in \mathcal{C}_{0, \infty, \mu_c}$,

$$\begin{aligned} \|g\|_{0, \infty, \mu_s, \nu} &\geq |g_s(\bar{x})| + \frac{|g_s(\bar{x}_1) - g_s(\bar{x}_2)|}{|\bar{x}_1 - \bar{x}_2|^\nu} \\ &\geq |f_s(\bar{x})| - |f_s(\bar{x}) - g_s(\bar{x})| \\ &\quad + \frac{|f_s(\bar{x}_1) - f_s(\bar{x}_2)|}{|\bar{x}_1 - \bar{x}_2|^\nu} - \frac{|(f_s(\bar{x}_1) - g_s(\bar{x}_1)) - (f_s(\bar{x}_2) - g_s(\bar{x}_2))|}{|\bar{x}_1 - \bar{x}_2|^\nu} \\ &= B + \delta - \left(|f_s(\bar{x}) - g_s(\bar{x})| + \frac{|(f_s(\bar{x}_1) - g_s(\bar{x}_1)) - (f_s(\bar{x}_2) - g_s(\bar{x}_2))|}{|\bar{x}_1 - \bar{x}_2|^\nu} \right) \\ &= B + \delta - \left(|f_c(\bar{x}) - g_c(\bar{x})| \frac{\mu_s(\bar{x})}{\mu_c(\bar{x})} + \frac{|(f_c(\bar{x}_1) - g_c(\bar{x}_1)) \frac{\mu_s(\bar{x}_1)}{\mu_c(\bar{x}_1)} - (f_c(\bar{x}_2) - g_c(\bar{x}_2)) \frac{\mu_s(\bar{x}_2)}{\mu_c(\bar{x}_2)}|}{|\bar{x}_1 - \bar{x}_2|^\nu} \right). \end{aligned}$$

For all $g \in \mathcal{C}_{0, \infty, \mu_c}$ with

$$\|f - g\|_{0, \infty, \mu_c} = \|f_c - g_c\|_{0, \infty} \leq \varepsilon$$

we have

$$|f_c(\bar{x}) - g_c(\bar{x})| \frac{\mu_s(\bar{x})}{\mu_c(\bar{x})} + \frac{|(f_c(\bar{x}_1) - g_c(\bar{x}_1)) \frac{\mu_s(\bar{x}_1)}{\mu_c(\bar{x}_1)} - (f_c(\bar{x}_2) - g_c(\bar{x}_2)) \frac{\mu_s(\bar{x}_2)}{\mu_c(\bar{x}_2)}|}{|\bar{x}_1 - \bar{x}_2|^\nu} \leq \varepsilon \frac{\mu_s(\bar{x})}{\mu_c(\bar{x})} + \frac{\varepsilon \frac{\mu_s(\bar{x}_1)}{\mu_c(\bar{x}_1)} + \varepsilon \frac{\mu_s(\bar{x}_2)}{\mu_c(\bar{x}_2)}}{|\bar{x}_1 - \bar{x}_2|^\nu}$$

by the triangle inequality. So suppose we choose ε small enough that the right hand side is $\leq \delta/2$.

Then for all $g \in \mathcal{C}_{0,\infty,\mu_c}$ with $\|f - g\|_{0,\infty,\mu_c} \leq \varepsilon$ we have

$$\begin{aligned} \|g\|_{0,\infty,\mu_s,\nu} &\geq B + \delta - \frac{\delta}{2} \\ &> B. \end{aligned}$$

Hence $g \in \Theta_0^c$ for all such g . Thus Θ_0^c is $\|\cdot\|_{0,m,\mu_c}$ -open and hence Θ_0 is $\|\cdot\|_{0,m,\mu_c}$ -closed.

Step 2 (Induction Step): This step follows the same arguments as those with bounded support. As in step 1, the main idea is simply to replace g with either g_c or g_s , as appropriate. \square

Proof of theorem 8 (Closedness for weighted norms on bounded domains). This proof is identical to the proof of theorem 6, except that now we use the compact embedding results of theorem 7 when necessary. \square

H Proofs of propositions from section 4

Proof of proposition 1. This proof is straightforward and we therefore omit it. \square

Proof of proposition 2. This proof is straightforward and we therefore omit it. \square

Proof of proposition 3. This proof is given in Gallant and Nychka (1987) as lemma A.2, and hence we omit it. \square

Proof of proposition 4. This proof is similar to the proof of proposition 3, which was shown in lemma A.2 of Gallant and Nychka (1987). Let $\mathcal{C} \subseteq \mathcal{D}$ be compact. We prove the proposition by induction on m (letting $m_0 = 0$, since it is irrelevant for the present result). For the base case, $m = 0$, the result holds trivially by letting $K_{\mathcal{C}} = 1$. Next suppose it holds for $m - 1$. Choose λ such that $|\lambda| = m$ and let $\nabla^\lambda = \nabla^\beta \nabla^\alpha$ where $|\alpha| = 1$ and $|\beta| = m - 1$. The result holds trivially if $\delta_s = 0$, so let $\delta_s \neq 0$. Then

$$\begin{aligned} \nabla^\lambda[\mu_s^{1/2}(x)] &= \nabla^\lambda \left[\exp \left(\frac{\delta_s}{2}(x'x) \right) \right] \\ &= \nabla^\beta \left(\nabla^\alpha \left[\exp \left(\frac{\delta_s}{2}(x'x) \right) \right] \right) \\ &= \nabla^\beta \left(\frac{\delta_s}{2} \exp \left(\frac{\delta_s}{2}(x'x) \right) \cdot \nabla^\alpha(x'x) \right) \\ &= \frac{\delta_s}{2} \sum_{\gamma \leq \beta} \begin{bmatrix} \beta \\ \gamma \end{bmatrix} \nabla^\gamma \left[\exp \left(\frac{\delta_s}{2}(x'x) \right) \right] \nabla^{\alpha+\beta-\gamma}(x'x) \\ &= \frac{\delta_s}{2} \sum_{\gamma \leq \beta} \begin{bmatrix} \beta \\ \gamma \end{bmatrix} [\nabla^\gamma \mu_s^{1/2}(x)] \nabla^{\alpha+\beta-\gamma}(x'x). \end{aligned}$$

In the fourth line we used Leibniz's formula. Next,

$$\begin{aligned} |\nabla^{\alpha+\beta-\gamma}(x'x)| &\leq \sum_{i=1}^{d_x} (x_i^2 + 2|x_i| + 2) \\ &\leq 4(1 + x'x). \end{aligned}$$

Hence

$$\begin{aligned} |\nabla^\lambda[\mu_s^{1/2}(x)]| &\leq \frac{|\delta_s|}{2} \sum_{\gamma \leq \beta} \begin{bmatrix} \beta \\ \gamma \end{bmatrix} |\nabla^\gamma \mu_s^{1/2}(x)| \cdot |4(1 + x'x)| \\ &\leq 2|\delta_s| \sum_{\gamma \leq \beta} \begin{bmatrix} \beta \\ \gamma \end{bmatrix} K_{\mathcal{C},m-1} \mu_s^{1/2}(x) \cdot |1 + x'x| \\ &\leq 2|\delta_s| \sum_{\gamma \leq \beta} \begin{bmatrix} \beta \\ \gamma \end{bmatrix} K_{\mathcal{C},m-1} \mu_s^{1/2}(x) \cdot M_{\mathcal{C}} \\ &= \mu_s^{1/2}(x) \left(2|\delta_s| \sum_{\gamma \leq \beta} \begin{bmatrix} \beta \\ \gamma \end{bmatrix} K_{\mathcal{C},m-1} \cdot M_{\mathcal{C}} \right). \end{aligned}$$

Here $M_{\mathcal{C}} = \sup_{x \in \mathcal{C}} |1 + x'x|$, which is finite since \mathcal{C} is compact. The second line follows by the induction hypothesis. \square

Proof of proposition 5. Pick $g(x) = 1 + x'x$. Notice that $g(x) \rightarrow \infty$ as $\|x\|_e \rightarrow \infty$. We prove the result by showing that for any $0 \leq |\lambda| \leq m_0$,

$$\nabla^\lambda \tilde{\mu}_c^{1/2}(x) = \exp \left[\frac{\delta_c}{2} (x'x) \right] \cdot p_\lambda(x) \quad (*)$$

for some polynomial $p_\lambda(x)$. Consequently, dividing by $\mu_s^{1/2}(x)$ yields

$$\frac{\nabla^\lambda \tilde{\mu}_c^{1/2}(x)}{\mu_s^{1/2}(x)} = \exp \left[\frac{\delta_c - \delta_s}{2} (x'x) \right] \cdot p_\lambda(x).$$

Since $\delta_c < \delta_s$,

$$\left| \frac{\nabla^\lambda \tilde{\mu}_c^{1/2}(x)}{\mu_s^{1/2}(x)} \right|$$

converges to zero as $\|x\|_e \rightarrow \infty$. This implies there is a J such that for all x with $\|x\|_e > J$, this ratio is smaller than M_1 . For all x with $\|x\|_e \leq J$, this ratio is a continuous function (the product of an exponential and a polynomial) on a compact set, and hence achieves a maximum M_2 . Let $M = \max\{M_1, M_2\}$. Thus the ratio is bounded by M for all $x \in \mathbb{R}^{d_x}$.

So it suffices to show equation (*). We proceed by induction. For the base case, $|\lambda| = 0$,

$$\begin{aligned}\nabla^0 \tilde{\mu}_c^{1/2}(x) &= \exp[\delta_c(x'x)/2] \cdot g(x) \\ &= \exp[\delta_c(x'x)/2] \cdot (1 + x^2).\end{aligned}$$

So the base case holds with $p_0(x) = g(x) = 1 + x^2$. Next, suppose it holds for $|\lambda| = m - 1$. Choose λ such that $|\lambda| = m$ and let $\nabla^\lambda = \nabla^\beta \nabla^\alpha$ where $|\alpha| = 1$ and $|\beta| = m - 1$. Then

$$\begin{aligned}\nabla^\lambda [\tilde{\mu}_c^{1/2}(x)] &= \nabla^\alpha [\nabla^\beta \tilde{\mu}_c^{1/2}(x)] \\ &= \nabla^\alpha [\exp[\delta_c(x'x)/2] \cdot p_\beta(x)] \\ &= \exp[\delta_c(x'x)/2] (\delta_c/2) p_\beta(x) \nabla^\alpha(x'x) + \exp[\delta_c(x'x)/2] \nabla^\alpha p_\beta(x) \\ &= \exp[\delta_c(x'x)/2] ((\delta_c/2) p_\beta(x) \nabla^\alpha(x'x) + \nabla^\alpha p_\beta(x)).\end{aligned}$$

Since the derivative of a polynomial is a polynomial, we're done. \square

Proof of proposition 6.

1. This follows immediately from lemmas 5 and 7:

$$\|\mu^{1/2} f\|_{m,2} \leq M_1 \|f\|_{m,2,\mu} \leq M_1 M \|\mu^{1/2} f\|_{m,2}.$$

2. This follows immediately from lemmas 6 and 8. \square

I Proofs of propositions from section 5

Proof of proposition 7. Suppose such a function μ existed. Define $g : (0, 1) \rightarrow \mathbb{R}$ by $g(x) = \log \mu(x)$. Then (1) implies that $g(x) \rightarrow -\infty$ as $x \rightarrow 0$. (2) implies that

$$g'(x) = \frac{1}{\mu(x)} \mu'(x) \leq K.$$

Hence $|g'(x)| \leq K$ for all $x \in (0, 1)$. This is a contradiction to $g(x) \rightarrow -\infty$ as $x \rightarrow 0$. \square

Proof of proposition 8. First consider the polynomial weight case, $\mu_s(x) = [x(1-x)]^{\delta_s}$. The proof is similar to the proof of propositions 3. We proceed by induction. For the base case $m = 0$, the result holds trivially by letting $K_C = 1$. Next suppose it holds for $m - 1$. If $\delta_s = 0$ the result holds

trivially, so let $\delta_s \neq 0$. We have

$$\begin{aligned}
\nabla^m[\mu_s^{1/2}(x)] &= \nabla^m \left([x(1-x)]^{\delta_s/2} \right) \\
&= \nabla^{m-1} \nabla^1 \left([x(1-x)]^{\delta_s/2} \right) \\
&= \nabla^{m-1} \left(\frac{\delta_s}{2} [x(1-x)]^{\delta_s/2-1} \nabla^1 [x(1-x)] \right) \\
&= \frac{\delta_s}{2} \sum_{\gamma \leq m-1} \begin{bmatrix} m-1 \\ \gamma \end{bmatrix} \nabla^\gamma \left([x(1-x)]^{\delta_s/2-1} \right) \nabla^{1+(m-1)-\gamma} [x(1-x)] \\
&= \frac{\delta_s}{2} \sum_{\gamma \leq m-1} \begin{bmatrix} m-1 \\ \gamma \end{bmatrix} \nabla^\gamma \left(\mu_{s,\tilde{\delta}}^{1/2}(x) \right) \nabla^{m-\gamma} [x(1-x)].
\end{aligned}$$

Here $\tilde{\delta} = \delta_s - 1/2$. $\nabla^n [x(1-x)]$ is either $x - x^2$ for $n = 0$, $1 - 2x$ for $n = 1$, -2 for $n = 2$, and 0 for $n > 2$. Hence

$$\begin{aligned}
M_{\mathcal{C}} &\equiv \sup_{x \in \mathcal{C}} |\nabla^{m-\gamma} [x(1-x)]| \\
&< \infty
\end{aligned}$$

since \mathcal{D} is bounded. So for all $x \in \mathcal{C}$,

$$\begin{aligned}
|\nabla^m[\mu_s^{1/2}(x)]| &\leq \frac{|\delta_s|}{2} \sum_{\gamma \leq m-1} \begin{bmatrix} m-1 \\ \gamma \end{bmatrix} |\nabla^\gamma[\mu_{s,\tilde{\delta}}^{1/2}(x)]| \cdot |\nabla^{m-\gamma}[x(1-x)]| \\
&\leq \frac{|\delta_s|}{2} \sum_{\gamma \leq m-1} \begin{bmatrix} m-1 \\ \gamma \end{bmatrix} K_{\mathcal{C},m-1} \mu_{s,\tilde{\delta}}^{1/2}(x) \cdot M_{\mathcal{C}} \\
&= \mu_{s,\tilde{\delta}}^{1/2}(x) \left(\frac{|\delta_s|}{2} \sum_{\gamma \leq m-1} \begin{bmatrix} m-1 \\ \gamma \end{bmatrix} K_{\mathcal{C},m-1} M_{\mathcal{C}} \right) \\
&= [x(1-x)]^{\delta_s/2-1} \left(\frac{|\delta_s|}{2} \sum_{\gamma \leq m-1} \begin{bmatrix} m-1 \\ \gamma \end{bmatrix} K_{\mathcal{C},m-1} M_{\mathcal{C}} \right) \\
&= \mu_s^{1/2}(x) \frac{1}{x(1-x)} \left(\frac{|\delta_s|}{2} \sum_{\gamma \leq m-1} \begin{bmatrix} m-1 \\ \gamma \end{bmatrix} K_{\mathcal{C},m-1} M_{\mathcal{C}} \right) \\
&\leq \mu_s^{1/2}(x) M'_{\mathcal{C}} \left(\frac{|\delta_s|}{2} \sum_{\gamma \leq m-1} \begin{bmatrix} m-1 \\ \gamma \end{bmatrix} K_{\mathcal{C},m-1} M_{\mathcal{C}} \right).
\end{aligned}$$

The second line follows by our $M_{\mathcal{C}}$ bound from above, and by the induction hypothesis with constant $K_{\mathcal{C},m-1}$. The last line follows since $\mathcal{C} \subseteq (0, 1)$ is compact, and hence x is bounded away from zero and one. So

$$M'_{\mathcal{C}} \equiv \sup_{x \in \mathcal{C}} \frac{1}{x(1-x)} < \infty.$$

Next consider the exponential weight case, $\mu_s(x) = \exp[\delta_s x^{-1}(1-x)^{-1}]$. The proof for this case is similar to the proofs of propositions 3 and 4. Let $\mathcal{C} \subseteq \mathcal{D}$ be compact. We prove the proposition by induction on m (letting $m_0 = 0$, since it is irrelevant for the present result). For the base case, $m = 0$, the result holds trivially by letting $K_{\mathcal{C}} = 1$. Next suppose it holds for $m - 1$. The result holds trivially if $\delta_s = 0$, so let $\delta_s \neq 0$. Then

$$\begin{aligned}
\nabla^m[\mu_s^{1/2}(x)] &= \nabla^m \left[\exp \left(\frac{\delta_s}{2} \frac{1}{x(1-x)} \right) \right] \\
&= \nabla^{m-1} \left(\nabla^1 \left[\exp \left(\frac{\delta_s}{2} \frac{1}{x(1-x)} \right) \right] \right) \\
&= \nabla^{m-1} \left(\frac{\delta_s}{2} \exp \left(\frac{\delta_s}{2} \frac{1}{x(1-x)} \right) \cdot \nabla^1 \left(\frac{1}{x(1-x)} \right) \right) \\
&= \frac{\delta_s}{2} \sum_{\gamma \leq m-1} \begin{bmatrix} m-1 \\ \gamma \end{bmatrix} \nabla^\gamma \left[\exp \left(\frac{\delta_s}{2} \frac{1}{x(1-x)} \right) \right] \nabla^{1+(m-1)-\gamma} \left(\frac{1}{x(1-x)} \right) \\
&= \frac{\delta_s}{2} \sum_{\gamma \leq m-1} \begin{bmatrix} m-1 \\ \gamma \end{bmatrix} [\nabla^\gamma \mu_s^{1/2}(x)] \nabla^{m-\gamma} \left(\frac{1}{x(1-x)} \right).
\end{aligned}$$

In the fourth line we used Leibniz's formula. Next, for any natural number n ,

$$\nabla^n \left(\frac{1}{x(1-x)} \right) = n! \sum_{j=0}^n \frac{(-1)^{n-j}}{(1-x)^{j+1} x^{n+1-j}}.$$

Hence

$$\begin{aligned}
|\nabla^m[\mu_s^{1/2}(x)]| &\leq \frac{|\delta_s|}{2} \sum_{\gamma \leq m-1} \begin{bmatrix} m-1 \\ \gamma \end{bmatrix} |\nabla^\gamma \mu_s^{1/2}(x)| \cdot \left| n! \sum_{j=0}^n \frac{(-1)^{n-j}}{(1-x)^{j+1} x^{n+1-j}} \right| \\
&\leq \frac{|\delta_s|}{2} \sum_{\gamma \leq m-1} \begin{bmatrix} m-1 \\ \gamma \end{bmatrix} |\nabla^\gamma \mu_s^{1/2}(x)| \cdot M_{\mathcal{C}} \\
&\leq \frac{|\delta_s|}{2} \sum_{\gamma \leq m-1} \begin{bmatrix} m-1 \\ \gamma \end{bmatrix} K_{\mathcal{C}, m-1} \mu_s^{1/2}(x) \cdot M_{\mathcal{C}} \\
&= \mu_s^{1/2}(x) \left(\frac{|\delta_s|}{2} \sum_{\gamma \leq m-1} \begin{bmatrix} m-1 \\ \gamma \end{bmatrix} K_{\mathcal{C}, m-1} \cdot M_{\mathcal{C}} \right).
\end{aligned}$$

Here

$$M_{\mathcal{C}} = \sup_{x \in \mathcal{C}} \left| n! \sum_{j=0}^n \frac{(-1)^{n-j}}{(1-x)^{j+1} x^{n+1-j}} \right|,$$

which is finite since $\mathcal{C} \subseteq (0, 1)$ is compact, and hence x is bounded away from zero and one. The third line follows by the induction hypothesis. \square

Proof of proposition 9. Let $g(x) = x^{-1}(1-x)^{-1}$. Then $g(x) \rightarrow \infty$ as $x \rightarrow 0$ or $x \rightarrow 1$. Note that

$\text{Bd}(\overline{\mathcal{D}}) = \{0, 1\}$. The rest of the proof is similar to that of proposition 5. It suffices to show that for any $0 \leq |\lambda| \leq m_0$,

$$\nabla^\lambda \tilde{\mu}_c^{1/2}(x) = \mu_c(x) \cdot r_\lambda(x) \quad (*)$$

for some rational function r_λ . Dividing (*) by $\mu_s^{1/2}(x)$ yields

$$\frac{\nabla^\lambda \tilde{\mu}_c^{1/2}(x)}{\mu_s^{1/2}(x)} = \exp[(\delta_c - \delta_s)g(x)] \cdot r_\lambda(x).$$

Since $\delta_c < \delta_s$, the absolute value of this expression converges to zero as $x \rightarrow 0$ or 1 . This proves part 2 of assumption 5. The proof of equation (*) is as in the proof of 5: The base case holds immediately with $r_0(x) = g(x)$. The induction step follows since the derivative of a rational function is still rational. \square

J Completeness of Sobolev spaces

When switching from the Sobolev sup-norm to the Sobolev L_p norm, a natural first space to consider is

$$\{f \in \mathcal{C}_m(\mathcal{D}) : \|f\|_{m,p,\mu} < \infty\}.$$

This space equipped with the norm $\|\cdot\|_{m,p,\mu}$ is unfortunately *not* $\|\cdot\|_{m,p,\mu}$ -complete. For unweighted spaces, $\mu(x) \equiv 1$, we can instead consider the *completion* of this space, denoted by $\mathcal{H}_{m,p,1}(\mathcal{D})$. An important result from functional analysis³ known as the ‘H=W theorem’ states that this completion equals the Sobolev space

$$\mathcal{W}_{m,p}(\mathcal{D}) = \{f \in \mathcal{W}_m(\mathcal{D}) : \|f\|_{m,p,1} < \infty\}.$$

Hence the way to complete the initial space is simply to allow for weakly differentiable functions, in addition to functions which are classically differentiable.

For weighted spaces, the H=W theorem does not necessarily hold; see Zhikov (1998).⁴ For this reason, we follow the literature by defining the weighted Sobolev space

$$\mathcal{W}_{m,p,\mu}(\mathcal{D}) = \{f \in \mathcal{W}_m(\mathcal{D}) : \|f\|_{m,p,\mu} < \infty\}.$$

As mentioned in section 2, this space is $\|\cdot\|_{m,p,\mu}$ -complete.

³See theorem 3.17 in Adams and Fournier (2003).

⁴Similar results sometimes obtain, however. For example, see Kufner and Opic (1984) remark 4.8 and also the discussion in Zhikov (1998). Also see remark 4.1 of Kufner and Opic (1984).

K Discussion of assumption 5

To get some intuition for assumption 5, consider the one dimensional case $d_x = 1$. In this case, we can usually take $m_0 = 1$, since $m_0 > d_x/2$ is then satisfied (see theorem 3 below). Then

$$\begin{aligned} \frac{|\nabla^0 \tilde{\mu}_c^{1/2}(x)|}{\mu_s^{1/2}(x)} &= \left| \frac{\nabla^0[\mu_c^{1/2}(x)g(x)]}{\mu_s^{1/2}(x)} \right| \\ &\leq \left(\frac{\mu_c(x)}{\mu_s(x)} \right)^{1/2} |g(x)| \end{aligned}$$

and

$$\begin{aligned} \frac{|\nabla^1 \tilde{\mu}_c^{1/2}(x)|}{\mu_s^{1/2}(x)} &= \left| \frac{\nabla^1[\mu_c^{1/2}(x)g(x)]}{\mu_s^{1/2}(x)} \right| \\ &= \left| \frac{\nabla^1 \mu_c^{1/2}(x)}{\mu_s^{1/2}(x)} g(x) + \frac{\mu_c^{1/2}(x)}{\mu_s^{1/2}(x)} \nabla^1 g(x) \right| \\ &\leq \frac{|\nabla^1 \mu_c^{1/2}(x)|}{\mu_s^{1/2}(x)} |g(x)| + \left(\frac{\mu_c(x)}{\mu_s(x)} \right)^{1/2} |\nabla^1 g(x)|. \end{aligned}$$

So when $d_x = 1$ with $m_0 = 1$, a sufficient condition for 5 is that there is a function g that diverges to infinity in the tails, but whose levels diverge slow enough that

$$|g(x)| = o\left(\left[\frac{\mu_c(x)}{\mu_s(x)}\right]^{-1/2}\right) \quad \text{and} \quad |\nabla^1 g(x)| = o\left(\left[\frac{|\nabla^1 \mu_c^{1/2}(x)|}{\mu_s^{1/2}(x)}\right]^{-1}\right)$$

and whose first derivative also satisfies

$$|\nabla^1 g(x)| = o\left(\left[\frac{\mu_c(x)}{\mu_s(x)}\right]^{-1/2}\right).$$

For further intuition, suppose assumption 3 held for μ_c . Then for all $x \in \mathbb{R}^{d_x}$ and any $0 \leq |\lambda| \leq m_0$,

$$\begin{aligned} |\nabla^\lambda \mu_c^{1/2}(x)| &\leq K \mu_c^{1/2}(x) \\ &= K \left(\frac{\mu_c(x)}{\mu_s(x)} \right)^{1/2} \mu_s^{1/2}(x) \end{aligned}$$

and hence

$$\frac{|\nabla^\lambda \mu_c^{1/2}(x)|}{\mu_s^{1/2}(x)} \leq K \left(\frac{\mu_c(x)}{\mu_s(x)} \right)^{1/2}$$

Now suppose assumption 1 holds. Then the right hand side converges to zero as $\|x\|_e \rightarrow \infty$. Thus, in this special case, a sufficient condition for assumption 5 is that $|g(x)|$ and its derivative $|\nabla^1 g(x)|$ do not diverge faster than $\sqrt{\mu_c(x)/\mu_s(x)}$ converges to zero.

L Closure of differentiable functions

The following lemma shows that the Sobolev sup-norm closure of a Sobolev sup-norm (with more derivatives) ball is a Hölder space with exponent 1. We assume $d_x = 1$ for notational simplicity, but the result can be extended to $d_x > 1$.

Lemma S2. Let \mathcal{D} be a convex open subset of \mathbb{R} . Let $m, m_0 \geq 0$ be integers. Define

$$\Theta_D = \{f \in \mathcal{C}_{m+m_0+1}(\mathcal{D}) : \|f\|_{m+m_0+1,\infty} \leq B\}$$

and

$$\Theta_L = \{f \in \mathcal{C}_{m+m_0}(\mathcal{D}) : \|f\|_{m+m_0,\infty,1,1} \leq B\}.$$

Let $\bar{\Theta}_D$ be the $\|\cdot\|_{m,\infty}$ -closure of Θ_D . Then $\bar{\Theta}_D = \Theta_L$.

Proof. We prove equality by showing that $\bar{\Theta}_D \subseteq \Theta_L$ and $\Theta_L \subseteq \bar{\Theta}_D$.

1. ($\bar{\Theta}_D \subseteq \Theta_L$). Let $f \in \bar{\Theta}_D$. We will show that $f \in \Theta_L$. By the definition of the $\|\cdot\|_{m,\infty}$ -closure, there exists a sequence $f_n \in \Theta_D$ such that

$$\|f_n - f\|_{m,\infty} \rightarrow 0.$$

Since $f_n \in \Theta_D$,

$$\|f_n\|_{m+m_0+1,\infty} = \max_{0 \leq |\lambda| \leq m+m_0+1} \sup_{x \in \mathcal{D}} |\nabla^\lambda f_n(x)| \leq B.$$

Also notice that for all $x, y \in \mathcal{D}$,

$$\frac{|\nabla^{m+m_0} f_n(x) - \nabla^{m+m_0} f_n(y)|}{|x - y|} \leq |\nabla^{m+m_0+1} f_n(\tilde{x})| \leq \sup_{x \in \mathcal{D}} |\nabla^{m+m_0+1} f_n(x)|$$

where \tilde{x} is between x and y , by the mean value theorem and convexity of \mathcal{D} . It follows that

$$\max_{|\lambda| \leq m+m_0} \sup_{x \in \mathcal{D}} |\nabla^\lambda f_n(x)| + \max_{|\lambda| = m+m_0} \sup_{x, y \in \mathcal{D}, x \neq y} \frac{|\nabla^\lambda f_n(x) - \nabla^\lambda f_n(y)|}{|x - y|} \leq \|f_n\|_{m+m_0+1,\infty} \leq B$$

and therefore $f_n \in \Theta_L$. But from part 5 of Theorem 2 we know that Θ_L is $\|\cdot\|_{m,\infty}$ -closed and since $\|f_n - f\|_{m,\infty} \rightarrow 0$ it follows that $f \in \Theta_L$.

2. ($\Theta_L \subseteq \bar{\Theta}_D$) Let $f \in \Theta_L$. We will show that $f \in \bar{\Theta}_D$. Specifically, we will show how to $\|\cdot\|_{m,\infty}$ -approximate f by a sequence of functions \tilde{f}_n in Θ_D . Define

$$M_1 = \max_{|\lambda| \leq m+m_0} \sup_{x, y \in \mathcal{D}, x \neq y} \frac{|\nabla^\lambda f(x) - \nabla^\lambda f(y)|}{|x - y|} < \infty$$

and

$$M_2 = \sup_{x, y \in \mathcal{D}, x \neq y} \frac{|\nabla^{m+m_0} f(x) - \nabla^{m+m_0} f(y)|}{|x - y|} < \infty.$$

If $\mathcal{D} \neq \mathbb{R}$, then since $\nabla^{m+m_0} f$ is Lipschitz, the Kirszbraun theorem (e.g., theorem 6.1.1 on page 189 of Dudley 2002) allows us to extend $\nabla^{m+m_0} f$ to a function “ $\nabla^{m+m_0} F$ ” on \mathbb{R} with the same Lipschitz constant. Define F to be the $m + m_0$ times antiderivative of $\nabla^{m+m_0} F$. Then F is $(m + m_0)$ -times differentiable, $\nabla^{m+m_0} F$ is Lipschitz with constant M_2 , and $F|_{\mathcal{D}} = f$. In particular, for this extension F ,

$$\max_{|\lambda| \leq m+m_0} \sup_{x,y \in \mathbb{R}, x \neq y} \frac{|\nabla^\lambda F(x) - \nabla^\lambda F(y)|}{|x - y|} = M_1$$

and

$$\sup_{x,y \in \mathbb{R}, x \neq y} \frac{|\nabla^{m+m_0} F(x) - \nabla^{m+m_0} F(y)|}{|x - y|} = M_2.$$

From here on we let $f(x) = F(x)$ denote the value of this extension of f if $x \notin \mathcal{D}$. The main issue is that f is only $(m+m_0)$ -times differentiable, but we want to approximate it by functions that are just a little bit smoother—functions that are $(m + m_0 + 1)$ -times differentiable. To do this, we convolve f with a smoother function:

$$f_n(x) = [f * \psi_{\varepsilon_n}](x) = \int_{\mathbb{R}} f(x + \varepsilon_n y) \psi(y) dy.$$

Here $*$ denotes convolution. ε_n is a sequence with $\varepsilon_n \rightarrow 0$ as $n \rightarrow \infty$. ψ_{ε_n} is an approximation to the identity: a function $\psi_{\varepsilon_n}(u) = \psi(u/\varepsilon_n)/\varepsilon_n$ where $\psi : \mathbb{R} \rightarrow \mathbb{R}$ is a $(m + m_0 + 1)$ -times continuously differentiable function such that $\psi(y) \geq 0$ for all $y \in \mathbb{R}$, $\psi(y) = 0$ if $|y| \geq 1$, and $\int_{-1}^1 \psi(y) dy = 1$. For example,

$$\psi(y) = B_k(1 - y^2)^k \mathbb{1}(|y| \leq 1).$$

where $k > m + m_0 + 1$ and B_k is such that the function integrates to 1. Note that f_n is $(m + m_0 + 1)$ -times differentiable.

For all $\lambda \leq m + m_0$,

$$\begin{aligned} [\nabla^\lambda f_n](x) &= [\nabla^\lambda f * \psi_{\varepsilon_n}](x) \\ &= \int_{\mathbb{R}} [\nabla^\lambda f](x - z) \frac{1}{\varepsilon_n} \psi\left(\frac{z}{\varepsilon_n}\right) dz \\ &= \int_{-1}^1 [\nabla^\lambda f](x - \varepsilon_n y) \psi(y) dy. \end{aligned}$$

The last line follows by a change of variables and since ψ is zero outside $[-1, 1]$. Hence

$$\begin{aligned}
|\nabla^\lambda f_n(x) - \nabla^\lambda f(x)| &\leq \int_{-1}^1 |\nabla^\lambda f(x - \varepsilon_n y) - \nabla^\lambda f(x)| \psi(y) dy \\
&\leq \int_{-1}^1 |M_1 \varepsilon_n y| \psi(y) dy \\
&= \varepsilon_n M_1 \int_{-1}^1 |y| \psi(y) dy \\
&\equiv \delta_n
\end{aligned}$$

for all $\lambda \leq m + m_0$. The first line follows since ψ integrates to 1. Since $\delta_n \rightarrow 0$, it follows that

$$\|f_n - f\|_{m+m_0, \infty} \rightarrow 0.$$

Moreover,

$$\begin{aligned}
|\nabla^{m+m_0} f_n(x_1) - \nabla^{m+m_0} f_n(x_2)| &\leq \int |\nabla^{m+m_0} f(x_1 - \varepsilon_n y) - \nabla^{m+m_0} f(x_2 - \varepsilon_n y)| \psi(y) dy \\
&\leq M_2 |x_1 - x_2|.
\end{aligned}$$

Since f_n is $(m + m_0 + 1)$ -times continuously differentiable,

$$|\nabla^{m+m_0+1} f_n(x)| = \lim_{h \rightarrow 0} \frac{|\nabla^{m+m_0} f_n(x+h) - \nabla^{m+m_0} f_n(x)|}{h} \leq M_2$$

for each $x \in \mathbb{R}$. Recall that

$$M_2 = \sup_{x, y \in \mathcal{D}, x \neq y} \frac{|\nabla^{m+m_0} f(x) - \nabla^{m+m_0} f(y)|}{|x - y|}.$$

This implies that

$$\begin{aligned}
\|f_n\|_{m+m_0+1, \infty} &\leq \|f_n\|_{m+m_0, \infty} + \sup_{x \in \mathcal{D}} |\nabla^{m+m_0+1} f_n(x)| \\
&\leq \|f\|_{m+m_0, \infty} + \|f_n - f\|_{m+m_0, \infty} + \sup_{x \in \mathcal{D}} |\nabla^{m+m_0+1} f_n(x)| \\
&\leq \|f\|_{m+m_0, \infty} + \delta_n + \sup_{x \in \mathcal{D}} |\nabla^{m+m_0+1} f_n(x)| \\
&\leq \left(\|f\|_{m+m_0, \infty} + \sup_{x, y \in \mathcal{D}, x \neq y} \frac{|\nabla^{m+m_0} f(x) - \nabla^{m+m_0} f(y)|}{|x - y|} \right) + \delta_n \\
&\leq B + \delta_n.
\end{aligned}$$

The last line follows since $f \in \Theta_L$. Thus f_n is *almost* in Θ_D , but not quite. But we can just

rescale f_n to put it inside Θ_D : Let

$$\tilde{f}_n(x) = \frac{B}{B + \delta_n} f_n(x).$$

Then $\|\tilde{f}_n\|_{m+m_0+1,\infty} \leq B$ and so $\tilde{f}_n \in \Theta_D$. Moreover,

$$\begin{aligned} \|\tilde{f}_n - f\|_{m,\infty} &\leq \|\tilde{f}_n - f\|_{m+m_0,\infty} \\ &\leq \|\tilde{f}_n - f_n\|_{m+m_0,\infty} + \|f_n - f\|_{m+m_0,\infty} \\ &= \max_{0 \leq |\lambda| \leq m+m_0} \sup_{x \in \mathcal{D}} \left| \nabla^\lambda \left(\frac{B}{B + \delta_n} f_n(x) \right) - \nabla^\lambda f_n(x) \right| + \|f_n - f\|_{m+m_0,\infty} \\ &= \left| \frac{B}{B + \delta_n} - 1 \right| \|f_n\|_{m+m_0,\infty} + \|f_n - f\|_{m+m_0,\infty} \\ &= \frac{\delta_n}{B + \delta_n} \|f_n\|_{m+m_0,\infty} + \|f_n - f\|_{m+m_0,\infty}. \end{aligned}$$

Since $\|f_n\|_{m+m_0,\infty} \leq \|f_n\|_{m+m_0+1,\infty} \leq B + \delta_n$,

$$\frac{\delta_n}{B + \delta_n} \|f_n\|_{m+m_0,\infty} \rightarrow 0.$$

We also know that $\|f_n - f\|_{m+m_0,\infty} \rightarrow 0$. It follows that

$$\|\tilde{f}_n - f\|_{m,\infty} \rightarrow 0.$$

But remember that $\tilde{f}_n \in \Theta_D$. So, by definition of the $\|\cdot\|_{m,\infty}$ -closure, $f \in \bar{\Theta}_D$.

□

M Sup-norm convergence over closed domains \mathcal{D}

Throughout the paper we have focused on functions with open domains \mathcal{D} . In practice we may also be interested in functions with closed domains \mathcal{D} . First, note that convergence of a sequence of functions in a Sobolev L_p norm where the integral is taken over the interior of \mathcal{D} implies convergence in the Sobolev L_p norm where the integral is taken over the entire \mathcal{D} . This follows since \mathcal{D} is a subset of \mathbb{R}^{d_x} and hence its boundary has measure zero. So the value of the integral is not affected by its values on the boundary. For Sobolev sup-norms, however, convergence over the interior of \mathcal{D} does not automatically imply convergence over all of \mathcal{D} . In the following lemma, we illustrate how to do this extension for sequences from a Hölder ball which are known to converge in the ordinary sup-norm over the interior. Similar results can be obtained with different parameter spaces and for convergence in general Sobolev sup-norms.

Lemma S3. Let $\mathcal{D} \subseteq \mathbb{R}^{d_x}$ be closed and convex. Let $f_n : \mathcal{D} \rightarrow \mathbb{R}$ be a sequence of functions in

$$\Theta = \{f \in \mathcal{C}_0(\mathcal{D}) : \|f\|_{0,\infty,1,\nu} \leq B\}.$$

Suppose

$$\sup_{x \in \text{int}\mathcal{D}} |f_n(x) - f(x)| \rightarrow 0.$$

for some function f . Suppose f is continuous at each boundary point in \mathcal{D} . Then

$$\sup_{x \in \mathcal{D}} |f_n(x) - f(x)| \rightarrow 0.$$

Proof of lemma S3. We want to show that for any $\varepsilon > 0$, there is an N such that

$$|f_n(x) - f(x)| \leq \varepsilon$$

for all $n \geq N$, for all $x \in \mathcal{D}$. For each $x \in \mathcal{D}$, choose an element $z_x \in \text{int}\mathcal{D}$ such that $\|x - z_x\|_e^\nu \leq \varepsilon/(3B)$ and

$$|f(x) - f(z_x)| \leq \frac{\varepsilon}{3}.$$

This is possible since f is continuous on all of \mathcal{D} , including at boundary points, and by convexity of \mathcal{D} . By the triangle inequality,

$$\begin{aligned} |f_n(x) - f(x)| &= |f_n(x) - f(x) - f_n(z_x) + f_n(z_x) - f(z_x) + f(z_x)| \\ &\leq |f_n(x) - f_n(z_x)| + |f(x) - f(z_x)| + |f_n(z_x) - f(z_x)|. \end{aligned}$$

By the definition of this parameter space we have

$$\sup_{x \in \mathcal{D}} |f_n(x) - f_n(z_x)| \leq B\|x - z_x\|_e^\nu \leq \frac{\varepsilon}{3}.$$

By uniform convergence of f_n to f on the interior of \mathcal{D} , there is an N such that

$$|f_n(z_x) - f(z_x)| \leq \frac{\varepsilon}{3}$$

for all $n \geq N$. Thus we're done. □

N Proofs for section 6

Proof of proposition 10. We omit this proof because it is almost identical to the proof of lemma A1 in Newey and Powell (2003). □

Proof of proposition 11. We verify the conditions of proposition 10.

1. The parameter space is $\|\cdot\|_c$ -compact by part 1 of theorems 3 and 4. Since the sieve space is a $\|\cdot\|_c$ -closed subset of the $\|\cdot\|_c$ -compact set Θ , it is also $\|\cdot\|_c$ -compact.

2. Define $Q(g) = -\mathbb{E}((Y - g(X))^2)$. Then for $g_1, g_2 \in \Theta$,

$$\begin{aligned}
& |Q(g_1) - Q(g_2)| \\
&= |\mathbb{E}(g_2(X)^2 - g_1(X)^2) + \mathbb{E}(2Y(g_1(X) - g_2(X)))| \\
&\leq |\mathbb{E}(g_2(X)^2 - g_1(X)^2)| + |\mathbb{E}(2Y(g_1(X) - g_2(X)))| \\
&= |\mathbb{E}(g_2(X) - g_1(X))(g_2(X) + g_1(X))| + 2|\mathbb{E}(Y(g_1(X) - g_2(X)))| \\
&\leq \sqrt{\mathbb{E}((g_2(X) - g_1(X))^2) \mathbb{E}((g_2(X) + g_1(X))^2)} + 2\sqrt{\mathbb{E}(Y^2) \mathbb{E}((g_1(X) - g_2(X))^2)} \\
&\leq \sqrt{\mathbb{E}((g_2(X) - g_1(X))^2) \mathbb{E}(2g_2(X)^2 + 2g_1(X)^2)} + 2\sqrt{\mathbb{E}(Y^2) \mathbb{E}((g_1(X) - g_2(X))^2)}.
\end{aligned}$$

The fourth line follows from the Cauchy-Schwarz inequality and the last line from $(a + b)^2 \leq 2a^2 + 2b^2$ for any $a, b \in \mathbb{R}$. Next,

$$\mathbb{E}((g_1(X) - g_2(X))^2) \leq \left(\sup_{x \in \mathbb{R}} |g_1(x) - g_2(x)| \mu_c(x) \right)^2 \mathbb{E}(\mu_c(X)^{-2}) = \|g_1 - g_2\|_c^2 \cdot \mathbb{E}(\mu_c(X)^{-2}).$$

Moreover, for all $g \in \Theta$,

$$\begin{aligned}
\mathbb{E}(g(X)^2) &= \mathbb{E}(g(X)^2 \mu_c(X)^2 \mu_c(X)^{-2}) \\
&\leq \|g\|_c^2 \cdot \mathbb{E}(\mu_c(X)^{-2}) \\
&\leq C^2 \|g\|_s^2 \cdot \mathbb{E}(\mu_c(X)^{-2}) \\
&\leq C^2 B^2 \mathbb{E}(\mu_c(X)^{-2}).
\end{aligned}$$

The third line follows since $\mathscr{H}_{1,2,\mu_s}$ is embedded in $\mathscr{C}_{0,\infty,\mu_c}$, by part 1 of theorem 3. Therefore

$$|Q(g_1) - Q(g_2)| \leq 2 \left(BC \mathbb{E}(\mu_c(X)^{-2}) + \sqrt{\mathbb{E}(Y^2) \mathbb{E}(\mu_c(X)^{-2})} \right) \|g_1 - g_2\|_c.$$

Since $\mathbb{E}(Y^2) < \infty$ and $\mathbb{E}(\mu_c(X)^{-2}) < \infty$, Q is $\|\cdot\|_c$ -continuous. Similarly, let $\widehat{Q}_n(g) = -\frac{1}{n} \sum_{i=1}^n (Y_i - g(X_i))^2$. Identical arguments imply that

$$|\widehat{Q}_n(g_1) - \widehat{Q}_n(g_2)| \leq 2 \left(BC \frac{1}{n} \sum_{i=1}^n \mu_c(X_i)^{-2} + \sqrt{\left(\frac{1}{n} \sum_{i=1}^n Y_i^2 \right) \left(\frac{1}{n} \sum_{i=1}^n \mu_c(X_i)^{-2} \right)} \right) \|g_1 - g_2\|_c.$$

Hence \widehat{Q} is $\|\cdot\|_c$ -continuous.

3. Suppose $Q(g) = Q(g_0)$. Then $\mathbb{E}((Y - g(X))^2) = \mathbb{E}((Y - g_0(X))^2)$, which implies that $g(X) = g_0(X)$ almost everywhere. If $g(\bar{x}) \neq g_0(\bar{x})$ for some \bar{x} , then $g(\bar{x}) \neq g_0(\bar{x})$ in a neighborhood of \bar{x} by continuity of g_0 , a contradiction. Hence $g(x) = g_0(x)$ for all $x \in \mathbb{R}$. Thus $\|g - g_0\|_c = \sup_{x \in \mathbb{R}} |g(x) - g_0(x)| \mu_c(x) = 0$. Moreover,

$$Q(g_0) = -\mathbb{E}((Y - g_0(X))^2) > -\mathbb{E}(2Y^2 + 2g_0(X)^2) > -\infty.$$

4. For any $g_k \in \Theta_k$

$$\|g_k - g_0\|_c \leq \sup_{|x| \leq M} |g_k(x) - g(x)| \sup_{|x| \leq M} \mu_c(x) + \sup_{|x| \geq M} |(g_k(x) - g(x))\mu_s(x)| \sup_{|x| \geq M} \frac{\mu_c(x)}{\mu_s(x)}.$$

Let $\varepsilon > 0$. Since g_k and g_0 are in Θ ,

$$\sup_{|x| \geq M} |(g_k(x) - g(x))\mu_s(x)| \leq \|g_k - g\|_s \leq 2B.$$

Thus, since μ_c and μ_s satisfy assumption 1, we can choose M such that

$$\sup_{|x| \geq M} |(g_k(x) - g(x))\mu_s(x)| \sup_{|x| \geq M} \frac{\mu_c(x)}{\mu_s(x)} \leq \frac{\varepsilon}{2}.$$

By assumption, for a fixed M , we can pick k large enough to make $\sup_{|x| \leq M} |g_k(x) - g(x)|$ arbitrarily small. By μ_c^2 satisfying the integrability assumption 6 and continuity of μ_c , $\sup_{|x| \leq M} \mu_c(x) < \infty$. Hence we can pick k large enough so that

$$\sup_{|x| \leq M} |g_k(x) - g(x)| \sup_{|x| \leq M} \mu_c(x) \leq \frac{\varepsilon}{2}.$$

Thus $\|g_k - g_0\|_c \leq \varepsilon$. Hence we have shown that $\|g_k - g_0\|_c \rightarrow 0$ as $k \rightarrow 0$.

5. For all $g \in \Theta_{k_n} \subseteq \Theta$,

$$(Y - g(X))^2 \leq 2Y^2 + g(X)^2 \leq 2Y^2 + 2B^2C^2\mu_c(X)^{-2}.$$

Since $\mathbb{E}(Y^2) < \infty$ and $\mathbb{E}(\mu_c(X)^{-2}) < \infty$ we have

$$\mathbb{E} \left(\sup_{g \in \Theta} (Y - g(X))^2 \right) < \infty.$$

This domination condition combined with $\|\cdot\|_c$ -compactness of Θ allows us to apply Jennrich's uniform law of large numbers to get

$$\sup_{g \in \Theta_{k_n}} |\widehat{Q}_n(g) - Q(g)| \xrightarrow{P} 0$$

as $n \rightarrow \infty$.

□

Proof of proposition 12. The proof is similar to the one of proposition 11 and verifies the conditions of proposition 10.

1. This step is identical to the corresponding step in the proof of proposition 11.

2. Define $Q(g) = -\mathbb{E}((Y - g(X))^2 \mu_c(X)^2)$. Then for $g_1, g_2 \in \Theta$,

$$\begin{aligned} |Q(g_1) - Q(g_2)| &= |\mathbb{E}((g_2(X)^2 - g_1(X)^2) \mu_c(X)^2) + \mathbb{E}(2Y(g_1(X) - g_2(X)) \mu_c(X)^2)| \\ &\leq \sqrt{\mathbb{E}((g_2(X) - g_1(X))^2 \mu_c(X)^2) \mathbb{E}((g_2(X) + g_1(X))^2 \mu_c(X)^2)} \\ &\quad + 2\sqrt{\mathbb{E}(Y^2 \mu_c(X)^2) \mathbb{E}((g_1(X) - g_2(X))^2 \mu_c(X)^2)} \\ &\leq \sqrt{\mathbb{E}((g_2(X) - g_1(X))^2 \mu_c(X)^2) \mathbb{E}(2g_2(X)^2 \mu_c(X)^2 + 2g_1(X)^2 \mu_c(X)^2)} \\ &\quad + 2\sqrt{\mathbb{E}(Y^2 \mu_c(X)^2) \mathbb{E}((g_1(X) - g_2(X))^2 \mu_c(X)^2)}. \end{aligned}$$

Next,

$$\mathbb{E}((g_1(X) - g_2(X))^2 \mu_c(X)^2) \leq \|g_1 - g_2\|_c^2.$$

Moreover, for all $g \in \Theta$,

$$\mathbb{E}(g(X)^2 \mu_c(X)^2) \leq B^2 M_5^2.$$

Therefore

$$|Q(g_1) - Q(g_2)| \leq 2 \left(BM_5 + \sqrt{\mathbb{E}(Y^2 \mu_c(X)^2)} \right) \|g_1 - g_2\|_c.$$

Since $\mathbb{E}(Y^2 \mu_c(X)^2) < \infty$, Q is continuous. Similarly, let $\widehat{Q}_n(g) = -\frac{1}{n} \sum_{i=1}^n (Y_i - g(X_i))^2 \mu_c(X_i)^2$. Identical arguments imply that

$$|\widehat{Q}_n(g_1) - \widehat{Q}_n(g_2)| \leq 2 \left(BM_5 + \sqrt{\frac{1}{n} \sum_{i=1}^n Y_i^2 \mu_c(X_i)^2} \right) \|g_1 - g_2\|_c.$$

Hence \widehat{Q} is continuous.

3. As before, $\mathbb{E}((Y - g(X))^2 \mu_c(X)^2) = \mathbb{E}((Y - g_0(X))^2 \mu_c(X)^2)$ implies $g(X) \mu_c(X) = g_0(X) \mu_c(X)$ almost everywhere. If $g(\bar{x}) \neq g_0(\bar{x})$ for some \bar{x} , then $g(\bar{x}) \neq g_0(\bar{x})$ in a neighborhood of \bar{x} by continuity of g_0 . Moreover if $\mu_c(\bar{x}) > 0$, then $\mu_c(x) > 0$ with positive probability in a neighborhood of \bar{x} , which contradicts that $g(X) \mu_c(X) = g_0(X) \mu_c(X)$ almost everywhere. Thus, $g(\bar{x}) \neq g_0(\bar{x})$ implies $\mu_c(\bar{x}) = 0$. Therefore $\|g - g_0\|_c = 0$. Moreover,

$$Q(g_0) = -\mathbb{E}((Y - g_0(X))^2 \mu_c(X)^2) > -\mathbb{E}(2Y^2 \mu_c(X)^2 + 2g_0(X)^2 \mu_c(X)^2) > -\infty.$$

4. This step is identical to the corresponding step in the proof of proposition 11.

5. For all $g \in \Theta_{k_n} \subseteq \Theta$,

$$(Y - g(X))^2 \mu_c(X)^2 \leq 2Y^2 \mu_c(X)^2 + 2g(X)^2 \mu_c(X)^2 \leq 2Y^2 \mu_c(X)^2 + 2B^2 M_5^2.$$

This combined with $\mathbb{E}(Y^2 \mu_c(X)^2) < \infty$ let us apply Jennrich's uniform law of large numbers, which gives

$$\sup_{\theta \in \Theta_{k_n}} |\widehat{Q}_n(\theta) - Q(\theta)| \xrightarrow{P} 0.$$

□

Proof of proposition 13. Let $g_{k_n} \in \tilde{\Theta}_{k_n}$ such that $\|g_{k_n} - g_0\|_c \rightarrow 0$. Then $\|g_{k_n}\|_c \leq \|g_0\|_c + 1$ for n large enough. Moreover, $\|g_0\|_c \leq C\|g_0\|_s < \infty$. From the proof of proposition 12 we know that

$$|Q(g_{k_n}) - Q(g_0)| \leq 2 \left(M_5(\|g_0\|_c + 1) + \sqrt{\mathbb{E}(Y^2 \mu_c(X)^2)} \right) \|g_{k_n} - g_0\|_c$$

and

$$|\hat{Q}_n(g_{k_n}) - \hat{Q}_n(g_0)| \leq 2 \left(M_5(\|g_0\|_c + 1) + \sqrt{\frac{1}{n} \sum_{i=1}^n Y_i^2 \mu_c(X_i)^2} \right) \|g_{k_n} - g_0\|_c.$$

Now write

$$\hat{Q}_n(g_{k_n}) - Q(g_{k_n}) = \left(\hat{Q}_n(g_{k_n}) - \hat{Q}_n(g_0) \right) + \left(\hat{Q}_n(g_0) - Q(g_0) \right) + \left(Q(g_0) - Q(g_{k_n}) \right).$$

$\hat{Q}_n(g_0) - Q(g_0) = O_p(1/\sqrt{n})$ by the central limit theorem, which applies since $\mathbb{E}((Y - g_0(X))^4) < \infty$ and μ_c is uniformly bounded above. Thus,

$$\hat{Q}_n(g_{k_n}) - Q(g_{k_n}) = O_p(\|g_{k_n} - g_0\|_c + 1/\sqrt{n}).$$

Since $\max\{1/\sqrt{n}, \|g_{k_n} - g_0\|_c\} = O(\lambda_n)$, lemma A.3 in Chen and Pouzo (2012) implies that for some $M_0 > 0$ it holds that $\|g_0\|_s \leq M_0$ and

$$\tilde{g}_w \in \{g \in \mathcal{W}_{1,2,\mu_s} : \|g\|_{1,2,\mu_s} \leq M_0\}$$

with probability arbitrarily close to 1 for all large n . Hence it suffices to prove that $\|\bar{g}_w - g_0\|_c \xrightarrow{P} 0$, where

$$\bar{g}_w(x) = \operatorname{argmax}_{g \in \tilde{\Theta}_{k_n}^{M_0}} - \left(\frac{1}{n} \sum_{i=1}^n (Y_i - g(X_i))^2 \mu_c(X_i)^2 + \lambda_n \|g\|_s \right)$$

and $\tilde{\Theta}_{k_n}^{M_0} = \{g \in \tilde{\Theta}_{k_n} : \|g\|_s \leq M_0\}$.

Consistency now follows from proposition 12 under two additional arguments:

1. First, $\sup_{g \in \tilde{\Theta}_{k_n}^{M_0}} \lambda_n \|g\|_s \leq \lambda_n M_0 \rightarrow 0$ and therefore the sample objective function (including the penalty) still converges to Q uniformly over $g \in \tilde{\Theta}_{k_n}^{M_0}$.
2. Second, since $\tilde{\Theta}_{k_n}^{M_0}$ is finite dimensional, for any $g_1, g_2 \in \tilde{\Theta}_{k_n}^{M_0}$ there exists $D > 0$ such that $\|g_1\|_s - \|g_2\|_s \leq D\|g_1\|_c - \|g_2\|_c \leq D\|g_1 - g_2\|_c$. Hence the sample objective function (including the penalty) is still continuous on $\tilde{\Theta}_{k_n}^{M_0}$.

All other assumptions of proposition 10 hold using the same arguments as those in the proof of proposition 12. Thus $\|\bar{g}_w - g_0\|_c \xrightarrow{P} 0$ and hence $\|\tilde{g}_w - g_0\|_c \xrightarrow{P} 0$. □

Proof of proposition 14. The proof is adapted from the proof of theorem 4.3 in Newey and Powell (2003). Again we verify the conditions of proposition 10.

1. This step is identical to the corresponding step in the proof of proposition 11.

2a. Define $Q(g) = -\mathbb{E}(\mathbb{E}(Y - g(X) | Z)^2)$. For $g_1, g_2 \in \Theta$,

$$\begin{aligned}
& |\mathbb{E}(Y - g_1(X) | Z)^2 - \mathbb{E}(Y - g_2(X) | Z)^2| \\
&= |\mathbb{E}(2Y | Z)\mathbb{E}(g_2(X) - g_1(X) | Z) + \mathbb{E}(g_2(X) - g_1(X) | Z)\mathbb{E}(g_2(X) + g_1(X) | Z)| \\
&\leq |\mathbb{E}(2Y + g_2(X) + g_1(X) | Z)| \cdot |\mathbb{E}(g_2(X) - g_1(X) | Z)| \\
&= |\mathbb{E}((2g_0(X) + g_2(X) + g_1(X))\mu_c(X)\mu_c(X)^{-1} | Z)| \cdot |\mathbb{E}((g_2(X) - g_1(X))\mu_c(X)\mu_c(X)^{-1} | Z)| \\
&\leq 4BM_5|\mathbb{E}(\mu_c(X)^{-1} | Z)| \cdot M_5\|g_1 - g_2\|_c \cdot |\mathbb{E}(\mu_c(X)^{-1} | Z)| \\
&= 4BM_5^2\mathbb{E}(\mu_c(X)^{-1} | Z)^2\|g_1 - g_2\|_c \\
&\leq 4BM_5^2\mathbb{E}(\mu_c(X)^{-2} | Z)\|g_1 - g_2\|_c.
\end{aligned}$$

The fourth line uses $\mathbb{E}(U | Z) = 0$ and the last uses Jensen's inequality. Therefore

$$\begin{aligned}
|Q(g_1) - Q(g_2)| &\leq \mathbb{E}(|\mathbb{E}(Y - g_1(X) | Z)^2 - \mathbb{E}(Y - g_2(X) | Z)^2|) \\
&\leq 4BM_5^2\mathbb{E}(\mu_c(X)^{-2})\|g_1 - g_2\|_c.
\end{aligned}$$

Hence, Q is continuous.

2b. Let

$$\Theta_{k_n} = \left\{ g \in \Theta : g = \sum_{j=1}^{k_n} b_j p_j(x) \text{ for some } b_1, \dots, b_{k_n} \in \mathbb{R} \right\}.$$

Define P_Z as the $n \times k_n$ matrix with (i, j) th element $p_j(X_i)$. Let $Q_Z = P_Z(P_Z'P_Z)^-P_Z'$ where $(P_Z'P_Z)^-$ denotes the Moore-Penrose generalized inverse of $(P_Z'P_Z)$. Let Y and $g(X)$ be the $n \times 1$ vectors with elements Y_i and $g(X_i)$, respectively. Define $\widehat{Q}_n(g) = -\frac{1}{n}\|Q_Z(Y - g(X))\|^2$. Then for $g_1, g_2 \in \Theta$,

$$\begin{aligned}
& |\widehat{Q}_n(g_1) - \widehat{Q}_n(g_2)| \\
&= \left| \frac{1}{n}\|Q_Z(Y - g_1(X))\|^2 - \frac{1}{n}\|Q_Z(Y - g_2(X))\|^2 \right| \\
&\leq \frac{1}{n}\|Q_Z(g_1(X) - g_2(X))\| \cdot \|Q_Z(2Y - g_1(X) - g_2(X))\| \\
&\leq \frac{1}{n}\|g_1(X) - g_2(X)\| \cdot \|2Y - g_1(X) - g_2(X)\| \\
&= \sqrt{\frac{1}{n} \sum_{i=1}^n (g_1(X_i) - g_2(X_i))^2 \mu_c(X_i)^2 \mu_c(X_i)^{-2}} \sqrt{\frac{1}{n} \sum_{i=1}^n (2Y_i - g_1(X_i) - g_2(X_i))^2} \\
&\leq \left(\sqrt{\frac{1}{n} \sum_{i=1}^n \mu_c(X_i)^{-2}} \sqrt{\frac{1}{n} \sum_{i=1}^n 4Y_i^2 + 4B^2M_5^2\mu_c(X_i)^{-2}} \right) \|g_1 - g_2\|_c.
\end{aligned}$$

The second line follows because, by the Cauchy-Schwarz inequality,

$$|(a'a) - (b'b)| = |(a-b)'(a+b)| \leq \sqrt{(a-b)'(a-b)}\sqrt{(a+b)'(a+b)}$$

for all $a, b \in \mathbb{R}^n$. The third line follows because Q_Z is idempotent and thus $\|Q_Z b\| \leq \|b\|$ for all $b \in \mathbb{R}^n$. Hence \widehat{Q}_n is continuous.

3. By completeness, $Q(g) = -\mathbb{E}(\mathbb{E}(Y - g(X) | Z)^2) = 0$ implies that $g(x) = g_0(x)$ almost everywhere. Identical arguments as those in the proof of proposition 11 then imply that $\|g - g_0\|_c = 0$, by continuity of g_0 . Moreover,

$$Q(g_0) = -\mathbb{E}(\mathbb{E}(U | Z)^2) = 0 > -\infty.$$

4. Assumption 4 of proposition 10 holds using identical arguments as those in the proof of proposition 11.
5. Assumption 5 of proposition 10 requires convergence of \widehat{Q}_n to Q uniformly over the sieve spaces. We show this by applying corollary 2.2 in Newey (1991). Θ is $\|\cdot\|_c$ -compact, which is Newey's assumption 1. Q is $\|\cdot\|_c$ -continuous, which is Newey's equicontinuity assumption. Next, define

$$B_n = \left(\sqrt{\frac{1}{n} \sum_{i=1}^n \mu_c(X_i)^{-2}} \sqrt{\frac{1}{n} \sum_{i=1}^n 4Y_i^2 + 4B^2 M_5^2 \mu_c(X_i)^{-2}} \right)$$

and recall that

$$|\widehat{Q}_n(g_1) - \widehat{Q}_n(g_2)| \leq B_n \|g_1 - g_2\|_c.$$

By Kolmogorov's strong law of large numbers and the existence of the relevant moments, $B_n = O_p(1)$. Hence Newey's assumption 3A holds. All that remains is to show Newey's assumption 2, pointwise convergence: $|\widehat{Q}(g) - Q(g)| = o_p(1)$ for all $g \in \Theta$. First write

$$\begin{aligned} |\widehat{Q}(g) - Q(g)| &= \frac{1}{n} \sum_{i=1}^n \mathbb{E}(Y - g(X) | Z = Z_i)^2 - \mathbb{E}(\mathbb{E}(Y - g(X) | Z)^2) \\ &\quad + \frac{1}{n} \sum_{i=1}^n \left(\widehat{\mathbb{E}}(Y - g(X) | Z = Z_i)^2 - \mathbb{E}(Y - g(X) | Z = Z_i)^2 \right), \end{aligned}$$

where $\widehat{\mathbb{E}}(Y - g(X) | Z = Z_i)$ is the series estimator of the conditional expectation evaluated

at Z_i . For the first part notice that $\mathbb{E}(Y - g(X) \mid Z = Z_i)^2$ is iid and

$$\begin{aligned} \mathbb{E}(\mathbb{E}(Y - g(X) \mid Z)^2) &\leq \mathbb{E}(\mathbb{E}((Y - g(X))^2 \mid Z)) \\ &\leq \mathbb{E}(2Y^2 + 2g(X)^2) \\ &\leq 2\mathbb{E}(Y^2) + 2\mathbb{E}(\mu_c(X)^{-1})\|g\|_c^2 \\ &< \infty. \end{aligned}$$

It follows from Kolmogorov's strong law of large numbers that

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E}(Y - g(X) \mid Z = Z_i)^2 - \mathbb{E}(\mathbb{E}(Y - g(X) \mid Z)^2) \xrightarrow{p} 0.$$

Next, following Newey (1991), define ρ as the $n \times 1$ vector containing $Y_i - g(X_i)$ and h as the $n \times 1$ vector containing $\mathbb{E}(Y - g(X) \mid Z = Z_i)$. Then

$$\left| \frac{1}{n} \sum_{i=1}^n \left(\widehat{E}(Y - g(X) \mid Z = Z_i)^2 - \mathbb{E}(Y - g(X) \mid Z = Z_i)^2 \right) \right| = \left| \|Q_Z \rho\|^2 - \|h\|^2 \right| / n.$$

Since for all $a, b \in \mathbb{R}^n$ it holds that $a'a - b'b = (a - b)'(a + b) + 2b'(a - b)$,

$$\left| \|Q_Z \rho\|^2 - \|h\|^2 \right| / n \leq (\|Q_Z \rho - h\|^2 + 2\|h\| \cdot \|Q_Z \rho - h\|) / n.$$

Since

$$\|h\|^2 / n = \frac{1}{n} \sum_{i=1}^n \mathbb{E}(Y - g(X) \mid Z = Z_i)^2,$$

the previous arguments imply that $\|h\|^2 / n = O_p(1)$. It therefore suffices to prove that $\|Q_Z \rho - h\|^2 / n = o_p(1)$, which by Markov's inequality is implied by

$$\mathbb{E}(\|Q_Z \rho - h\|^2) / n \rightarrow 0.$$

as $n \rightarrow \infty$. Newey (1991) shows

$$\mathbb{E}(\|Q_Z \rho - h\|^2) / n \leq \mathbb{E}(\text{trace}(Q_Z \text{var}(h \mid Z))) / n + o(1).$$

Therefore,

$$\begin{aligned}
\mathbb{E} (\|Q_Z \rho - h\|^2) / n &\leq \mathbb{E} \left(\sum_{i=1}^n (Q_Z)_{ii} \text{var}(Y_i - g(X_i) \mid Z_i) \right) / n + o(1) \\
&\leq \mathbb{E} \left(\sqrt{\frac{1}{n} \sum_{i=1}^n (Q_Z)_{ii}^2 \frac{1}{n} \sum_{i=1}^n \text{var}(Y_i - g(X_i) \mid Z_i)^2} \right) + o(1) \\
&\leq \mathbb{E} \left(\sqrt{\frac{1}{n} \text{trace}(Q_Z' Q_Z) \frac{1}{n} \sum_{i=1}^n \text{var}(Y_i - g(X_i) \mid Z_i)^2} \right) + o(1) \\
&= \mathbb{E} \left(\sqrt{\frac{1}{n} \text{trace}(Q_Z) \frac{1}{n} \sum_{i=1}^n \text{var}(Y_i - g(X_i) \mid Z_i)^2} \right) + o(1) \\
&\leq \sqrt{\frac{k_n}{n}} \mathbb{E} \left(\sqrt{\frac{1}{n} \sum_{i=1}^n \text{var}(Y_i - g(X_i) \mid Z_i)^2} \right) + o(1) \\
&\leq \sqrt{\frac{k_n}{n}} \sqrt{\mathbb{E} (\text{var}(Y_i - g(X_i) \mid Z_i)^2)} + o(1).
\end{aligned}$$

The second line follows from the Cauchy-Schwarz inequality. The third line from the definition of the trace. The fourth line because Q_Z is idempotent. The fifth line because $\text{trace}(Q_Z) \leq k_n$. The last line by Jensen's inequality. Since $\mathbb{E} \left((\text{var}(Y_i - g(X_i) \mid Z_i))^2 \right) < \infty$ and $k_n/n \rightarrow 0$, it follows that

$$\mathbb{E} (\|Q_Z \rho - h\|^2) / n \rightarrow 0.$$

□

References

- ADAMS, R. A. AND J. J. FOURNIER (2003): *Sobolev spaces*, vol. 140, Academic press, 2nd ed.
- BROWN, R. AND B. OPIC (1992): "Embeddings of weighted Sobolev spaces into spaces of continuous functions," *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 439, 279–296.
- CHEN, X. AND D. POUZO (2012): "Estimation of nonparametric conditional moment models with possibly nonsmooth generalized residuals," *Econometrica*, 80, 277–321.
- CONWAY, J. B. (1985): *A course in functional analysis*, Springer.
- DUDLEY, R. M. (2002): *Real analysis and probability*, vol. 74, Cambridge University Press.
- FOLLAND, G. B. (1999): *Real analysis: Modern techniques and their applications*, John Wiley & Sons, Inc., 2nd ed.
- GALLANT, A. AND D. NYCHKA (1987): "Semi-nonparametric maximum likelihood estimation," *Econometrica*, 55, 363–390.

- KUFNER, A. (1980): *Weighted Sobolev spaces*, BSB B. G. Teubner Verlagsgesellschaft.
- KUFNER, A. AND B. OPIC (1984): “How to define reasonably weighted Sobolev spaces,” *Commentationes Mathematicae Universitatis Carolinae*, 25, 537–554.
- NEWBY, W. K. (1991): “Uniform convergence in probability and stochastic equicontinuity,” *Econometrica*, 1161–1167.
- NEWBY, W. K. AND J. L. POWELL (2003): “Instrumental variable estimation of nonparametric models,” *Econometrica*, 71, 1565–1578.
- RODRÍGUEZ, J. M., V. ÁLVAREZ, E. ROMERA, AND D. PESTANA (2004): “Generalized weighted Sobolev spaces and applications to Sobolev orthogonal polynomials I,” *Acta Applicandae Mathematicae*, 80, 273–308.
- SANTOS, A. (2012): “Inference in nonparametric instrumental variables with partial identification,” *Econometrica*, 80, 213–275.
- TAO, T. (2010): *An epsilon of room, 1: Real analysis*, American Mathematical Society.
- ZHIKOV, V. V. (1998): “Weighted Sobolev spaces,” *Sbornik: Mathematics*, 189, 1139–1170.